arXiv:math/0401225v3 [math.AG] 14 Sep 2004

# DANIELEWSKI-FIESELER SURFACES

ADRIEN DUBOULOZ

Abstract. We study a class of normal affine surfaces with additive group actions which contains in particular the Danielewski surfaces in $\mathbb{A}^3$ given by the equations $x^n z = P(y)$, where $P$ is a nonconstant polynomial with simple roots. We call them Danielewski-Fieseler Surfaces. We reinterpret a construction of Fieseler [7] to show that these surfaces appear as the total spaces of certain torsors under a line bundle over a curve with an $r$-fold point. We classify Danielewski-Fieseler surfaces through labelled rooted trees attached to such a surface in a canonical way. Finally, we characterize those surfaces which have a trivial Makar-Limanov invariant in terms of the associated trees.

## Introduction

The surfaces $S_j = \left\{x^j z = y^2 + 1\right\}$, $j \geq 1$, in $\mathbb{C}^3 = \operatorname{Spec}(\mathbb{C}[x, y, z])$ admit free actions of the additive group $\mathbb{G}_{a,\mathbb{C}}$ induced by the locally nilpotent derivations $\partial_j = x^j \partial_y + 2y\partial_z$ of $\mathbb{C}[x, y, z]$ respectively. Danielewski [4] observed that the associated quotient morphism $\pi : S_j \rightarrow S_j//\mathbb{G}_{a,\mathbb{C}} \simeq \operatorname{Spec}(\mathbb{C}[x])$ is an $\mathbb{A}^1$-fibration which factors throught an $\mathbb{A}^1$-bundle $\rho : S_j \rightarrow X$ over the affine line with a double origin $\delta : X \rightarrow \operatorname{Spec}(\mathbb{C}[x])$. Indeed, $S_j$ is obtained as the gluing of two copies $\operatorname{Spec}(\mathbb{C}[x][T_i])$ of $\mathbb{C}^2$, $i = 1, 2$, over $\mathbb{C}^* \times \mathbb{C}$ by means of the $\mathbb{C}[x]_x$-algebras isomorphisms $\mathbb{C}[x]_x[T_1] \xrightarrow{\sim} \mathbb{C}[x]_x[T_2]$, $T_1 \mapsto 2x^{-j} + T_2$. This interpretation was further generalized by Fieseler [7] to describe certain invariant neighbourhoods of the fibers of a quotient $\mathbb{A}^1$-fibration $\pi : S \rightarrow Z$ associated with a nontrivial action of $\mathbb{G}_{a,\mathbb{C}}$ on a normal affine surface $S$. More precisely, he established that if a fiber $\pi^{-1}(z_0)$ is reduced then the induced morphism $p_2 : S \times_Z \operatorname{Spec}(\mathcal{O}_{Z,z_0}) \rightarrow \operatorname{Spec}(\mathcal{O}_{Z,z_0})$, where $\mathcal{O}_{Z,z_0}$ denotes the local ring of $z_0 \in Z$, factors through an $\mathbb{A}^1$-bundle $\rho : S \rightarrow X$ over the scheme $X$ obtained from $\operatorname{Spec}(\mathcal{O}_{Z,z_0})$ by replacing $z_0$ by as many points as there are connected components in $\pi^{-1}(z_0)$. More generally, given a field $k$ of caracteristic zero and a pair $(X = \operatorname{Spec}(A), x_0 = \operatorname{div}(x))$, where $A$ is either discrete valuation ring with uniformizing parameter $x$ and residue field $k$ or of a polynomial ring $A = k[x]$, the same description holds for the following class of surfaces.

**Definition 0.1.** A *Danielewski-Fieseler surface with base* $(A, x)$ (A DFS, for short) is an integral affine $X$-scheme $\pi : S \rightarrow X$ of finite type, restricting to a trivial line bundle over $X_* = X \setminus \{x_0\}$, and such that $\pi^{-1}(x_0)$ is nonempty and reduced, consisting of a disjoint union of curves isomorphic to affine line $\mathbb{A}^1_k$.

In this paper, we give a combinatorial description of DFS's in terms of $(A, x)$-*labelled rooted trees*, that is, pairs $\gamma = (\Gamma, \sigma)$ consisting of a rooted tree $\Gamma$ and a cochain $\sigma \in A^n$, indexed by the terminal elements of $\Gamma$, satisfying certain conditions with respect to the geometry of $\Gamma$ (see (1.5) below). Then, as an application, we

---

2000 *Mathematics Subject Classification.* 14L30,14R05,14R20,14R25.

characterize the DFS's with base $(k[x], x)$ which have a trivial Makar-Limanov invariant [10].

The paper is divided as follows. In section 1, we collect some preliminary results on labelled rooted trees. In section 2, we reinterpret the 'cocycle construction' of Fieseler [7] to describe DFS's as torsors under certain line bundles on a nonseparated scheme. In section 3, we associate to every labelled tree $\gamma$, a DFS $\pi_\gamma : S(\gamma) \to X$ which comes equipped with a canonical birational $X$-morphism $\psi_\gamma : S(\gamma) \to \mathbb{A}^1_X$. For instance, given a polynomial $P \in k[y]$ with simple roots $y_1, \ldots, y_n \in k$, the Danielewski-Fieseler surface with base $(k[x], x)$

$$\pi : S_{P,m} = \operatorname{Spec}\left(k[x,y,z] / (x^m z - P(y))\right) \longrightarrow X = \operatorname{Spec}(k[x]),$$

equipped with the morphism $pr_y : S_{P,m} \to \mathbb{A}^1_X = \operatorname{Spec}(k[x][y])$ corresponds to the following labelled tree $\gamma_{P,m} = (\Gamma_{m,n}, \sigma)$.

$\Gamma_{m,n} =$



$\sigma = \{y_i\}_{i=1,\ldots,n} \in k[x]^n$

In Theorem (3.2), we establish that the category $\overrightarrow{\mathfrak{D}}_{(A,x)}$ of DFS's with base $(A, x)$ equipped with certain birational morphisms as above and the category $\mathcal{T}_{(A,x)}$ of $(A, x)$-labelled trees are equivalent. Then, in Theorem (3.10), we classify these DFS's up to $X$-isomorphisms in terms of the corresponding trees. In section 4, we decompose birational $X$-morphisms between two DFS's into a succession of simple affine modifications [11], which we call *fibered modifications* (Theorem 4.5). This leads to a canonical procedure for constructing embeddings of a DFS $\pi : S \to X$ into a projective $X$-scheme $\bar{\pi} : \bar{S} \to X$ (Proposition 4.7). Section 5 is devoted to the study of DFS's with base $(k[x], x)$, which we call simply *Danielewski surfaces.* We recall that the Makar-Limanov invariant [10] of an affine variety $V/k$ is defined as the intersection in $\Gamma(V, \mathcal{O}_V)$ of all the invariant rings of $\mathbb{G}_{a,k}$-actions on $V$. Makar-Limanov [12] and [13] established that a surface $S_{P,m} \simeq S(\gamma_{P,m})$ as above has trivial Makar-Limanov invariant $\operatorname{ML}(S_{P,m}) = k$ if and only if $m = 1$ and $\deg(P) \geq 1$. More generally, in case that $k = \bar{k}$ is algebraically closed, we give the following characterization (Theorem 5.4).

**Theorem 0.2.** *A Danielewski surface $S(\gamma)$ has a trivial Makar-Limanov invariant if and only if $\gamma$ is a comb,* i.e *a labelled tree $\gamma = (\Gamma, \sigma)$ with the property that all but at most one of the direct descendants of a given $e \in \Gamma$ are terminal elements of $\Gamma$.*

Finally, we obtain the following description of normal affine surfaces with a trivial Makar-Limanov invariant, which generalizes previous results obtained by Daigle-Russell [3] and Miyanishi-Masuda [14] for the particular case of log $\mathbb{Q}$-homology planes (Theorem 5.9).

**Theorem 0.3.** *Every normal affine surface $S/\bar{k}$ with a trivial Makar-Limanov invariant is isomorphic to a cyclic quotient of a Danielewski surface.*

## 1. Preliminaries on rooted trees

Here we give some results on rooted trees and labelled rooted trees which will be used in the following sections.

**Basic facts on rooted trees.** Let $(G, \leq)$ be a nonempty, finite, partially ordered set (a *poset*, for short). A totally ordered subset $C \subset G$ is called a *subchain of length* $l(C) = \mathrm{Card}(C) - 1$. *A* subchain of maximal length is called a *maximal subchain.* For every $e \in G$, we let $(\uparrow e)_G = \{e' \in G, e' \geq e\}$ and $(\downarrow e)_G = \{e' \in G, e' \leq e\}$. The *edges* $E(G)$ of $G$ are the subsets $\overrightarrow{ee'} = \{e' < e\}$ of $G$ with two elements such that $(\uparrow e')_G \cap (\downarrow e)_G = \overrightarrow{ee'}$.

**Definition 1.1.** A *(rooted) tree* $\Gamma$ is poset with a unique minimal element $e_0$ called the *root*, and such that $(\downarrow e)_\Gamma$ is a chain for every $e \in \Gamma$. A *(rooted) subtree* of $\Gamma$ is a sub-poset $\Gamma' \subset \Gamma$ which is a rooted tree for the induced order. We say that $\Gamma'$ is *maximal* if there exists $e' \in \Gamma$ such that $\Gamma' = (\uparrow e')_\Gamma$.

**1.2.** An element $e \in \Gamma$ such that $l(\downarrow e)_\Gamma = m$ is said to be *at level* $m$. The maximal elements $e_i = e_{i,m_i}$, where $m_i = l(\downarrow e_i)_\Gamma$, of $\Gamma$ are called the *leaves* of $\Gamma$. We denote the set of those elements by $L(\Gamma)$. The corresponding subchains

$$(1)\qquad (\downarrow e_{i,m_i})_\Gamma = \{e_{i,0} = e_0 < e_{i,1} < \cdots < e_{i,m_i-1} < e_{i,m_i} = e_i\}, \quad i = 1, \ldots, n.$$

are the maximal subchains of $\Gamma$. We say that $\Gamma$ has *height* $h(\Gamma) = \max(m_i)$. The *first common ancestor* of two element $e, e' \in \Gamma \setminus \{e_0\}$ is the maximal element of the chain $((\downarrow e)_\Gamma \setminus \{e\}) \cap ((\downarrow e')_\Gamma \setminus \{e'\})$. The *children* of $e \in \Gamma \setminus L(\Gamma)$ are the minimal elements of $(\uparrow e)_\Gamma \setminus \{e\}$. We denote the set of those elements by $\mathrm{Ch}_\Gamma(e)$.



Figure 1. A tree $\Gamma$ rooted in $e_0$.

**Definition 1.3.** A *morphism of (rooted) trees* is an order-preserving map $\tau : \Gamma' \to \Gamma$ satisfying the following properties:
a) The image of a maximal chain of $\Gamma'$ by $\tau$ is a maximal chain of $\Gamma$.
b) For every $e' \in \Gamma'$, $\tau^{-1}(\tau(e'))$ is either $e'$ itself or a maximal subtree of $\Gamma'$.
Injective and bijective morphisms are referred to as embeddings and isomorphisms respectively.

**1.4.** This definition implies that for every leaf $e'_{i,m'_i}$ of $\Gamma'$ at level $m'_i$, $\tau\left(e'_{i,m'_i}\right)$ is a leaf $e_{j(i),m_{j(i)}}$ of $\Gamma$ at level $m_{j(i)} \leq m'_i$, such that $\tau\left(e'_{i,k}\right) = e_{j(i),\min\left(k,m_{j(i)}\right)}$ for every $k = 0, \ldots, m'_i$ (see (1) for the notation).

**Labelled rooted trees.** In this subsection, we fix a pair $(A,x)$ consisting of a domain $A$ and an element $x \in A \setminus \{0\}$.

**Definition 1.5.** An $(A,x)$-*labelled tree* is a pair $\gamma = (\Gamma, \sigma)$ consisting of a a tree $\Gamma$ with the leaves $e_{1,m_1}, \ldots, e_{n,m_n}$ and a cochain $\sigma = \{\sigma_i\}_{i=1,\ldots,n} \in A^n$ such that $\sigma_j - \sigma_i \in x^{d_{ij}} A \setminus x^{d_{ij}+1} A$ whenever the first common ancestor of $e_{i,m_i}$ and $e_{j,m_j}$ is at level $d_{ij} < \min(m_i, m_j)$. A cochain $\sigma$ with this this property is said to be $\Gamma$-*compatible*.

A *morphism of* $(A,x)$-*labelled trees* $\tau : (\Gamma', \sigma') \to (\Gamma, \sigma)$ is a morphism of trees $\tau : \Gamma' \to \Gamma$ such that if $e_{j(i),m_{j(i)}} \in L(\Gamma)$ is the image of a leaf $e'_{i,m'_i} \in L(\Gamma')$ by $\tau$, then $\sigma_i - \sigma_{j(i)} \in x^{m_{j(i)}} A$.

The category of $(A,x)$-labelled just defined is denoted by $\mathcal{T}_{(A,x)}$.

**Example 1.6.** Every isomorphism class of $(k[x], x)$-labelled trees contains a tree $\gamma = (\Gamma, \sigma)$ with the leaves $e_{1,m_1}, \ldots, e_{n,m_n}$, such that $\sigma_i = \sum_{j=0}^{m_i-1} w_{i,j} x^j \in k[x]$ is a polynomial of degree $< m_i$ for every $i = 1, \ldots, n$. In turn, this cochain $\sigma$ is uniquely determined by the choice of a *weight function*

$$w_\sigma : E(\Gamma) \longrightarrow k, \quad \overrightarrow{e_{i,j+1}e_{i,j}} \mapsto w_\sigma\left(\overrightarrow{e_{i,j+1}e_{i,j}}\right) = w_{i,j}$$

with the property that $w_\sigma\left(\overrightarrow{e'e}\right) \neq w_\sigma\left(\overrightarrow{e''e}\right)$ whenever $e'$ and $e''$ are children of the same $e \in \Gamma$. A tree $\Gamma$ equipped with such a function $w$ is referred to as a *fine $k$-weighted tree.* A morphism of trees $\tau : \Gamma' \to \Gamma$ gives rise to a morphism of $(k[x], x)$-labelled $\tau : (\Gamma', \sigma') \to (\Gamma, \sigma)$ if and only if $w_\sigma\left(\overrightarrow{\tau(e)\,\tau(e')}\right) = w_{\sigma'}\left(\overrightarrow{ee'}\right)$whenever $\tau(e) \neq \tau(e')$. If it is the case, then we say that $\tau$ is a *morphism of fine $k$-weighted trees.* In this way, we obtain a bijection between morphisms of $(A,x)$-labelled trees and morphisms of fine $k$-weighted trees. We conclude that the categories $\mathcal{T}_{(k[x],x)}$ of $(k[x], x)$-labelled trees and $\mathcal{T}^k_w$ of fine $k$-weighted trees are equivalent.

**Gluing trees.** Intuitively, $(A,x)$-labelled trees are constructed by gluing $(A,x)$-labelled chains. Similarly, a morphism $\tau : \gamma' \to \gamma$ of $(A,x)$-labelled trees is uniquely determined by its values on maximal subchains of $\gamma'$. More precisely, we have the following results.

**Proposition 1.7.** *Suppose that $(n, m, d, \sigma)$ is a data consisting of an integer $n \geq 1$, a multi-index $m = (m_1, \ldots, m_n) \in \mathbb{Z}^n_{>0}$, a matrix $d = (d_{ij})_{i,j=1,\ldots,n} \in Mat_n(\mathbb{Z}_{\geq 0})$ and a cochain $\sigma = \{\sigma_i\}_{i=1,\ldots,n} \in A^n$ with the following properties.*

*1) For every $i \neq j$ , $d_{ij} = d_{ji} < \min(m_i, m_j)$ .*

*2) For every triple $i, j, k$, $\min(d_{ij}, d_{ik}) = \min(d_{ji}, d_{jk})$.*

*3) For every $i \neq j$, $\sigma_j - \sigma_i \in x^{d_{ij}} A \setminus x^{d_{ij}+1} A$.*

*Then there exists a tree $\Gamma$, unique up to a unique isomorphism, with the root $e_0$ and the leaves $e_i$ at levels $m_i$, $i = 1, \ldots, n$, such that $\sigma$ is $\Gamma$-compatible.*

*Proof.* Up to an isomorphism, $\gamma_i = (C_i = \{e_{i,0} < e_{i,1} < \cdots < e_{i,m_i}\}, \sigma_i)$ is the unique chain of height $m_i \geq 1$ such that $\sigma_i$ is $C_i$-compatible. For every $i \neq j$, we let $C_{ij} = \left(\downarrow e_{i,d_{ij}}\right)_{C_i} \subset C_i$. Condition (2) guarantees that there exist isomorphisms of chains $\phi_{ij} : C_{ij} \overset{\sim}{\to} C_{ij}$ such that $\phi_{ij}(C_{ij} \cap C_{ik}) = C_{ji} \cap C_{jk}$ for every triple $i, j, k$, and such that the cocycle condition $\phi_{ik} = \phi_{jk} \circ \phi_{ij}$ holds on on $C_{ij} \cap C_{ik}$. Moreover, (1) implies that for every $i = 1, \ldots, n$, $e_{i,0}$ belongs to $C_{ij}$ whereas $e_{i,m_i}$ does not. Therefore, the quotient poset $\Gamma = \bigsqcup_{i=1}^n C_i / (C_{ij} \ni e_{i,k} \sim \phi_{ij}(e_{j,k}) \in C_{ji})$ is a tree rooted in the common image $e_0 = \overline{e_{i,0}}$ of the roots $e_{i,0}$ of the chains $C_i$, and with the leaves $e_i = \overline{e_{i,m_i}}$ at level $m_i$, $i = 1, \ldots, n$ . Finally, (3) means exactly that $\sigma$ is $\Gamma$-compatible. This completes the proof. $\square$

**Proposition 1.8.** *Let $\gamma = (\Gamma, \sigma)$ and $\gamma' = (\Gamma', \sigma')$ be two $(A, x)$-labelled trees. Suppose we are given a collection of morphisms of $(A, x)$-labelled trees*

$$\tau_i : \gamma'_i = (\Gamma'_i, \sigma'_i) = ((\downarrow e'_i)_{\Gamma'}, \sigma'_i) \longrightarrow \gamma, \quad \textit{where } L(\Gamma') = \{e'_1, \ldots, e'_{n'}\},$$

*restricting to a same morphism of trees $\tau_{ij} = \tau_{ji} : \Gamma'_i \cap \Gamma'_j \to \tau_i(\Gamma'_i) \cap \tau_j\left(\Gamma'_j\right)$ for every $i \neq j$. Then there exists a unique morphism of $(A, x)$-labelled trees $\tau : \gamma' \to \gamma$ such that $\tau \mid_{\gamma'_i} = \tau_i$ for every $i = 1, \ldots, n'$.*

*Proof.* The conditions guarantee that there exists a unique order-preserving map $\tau : \Gamma' \to \Gamma$ such that $\tau \mid_{\Gamma'_i} = \tau_i$, and such that the preimage $\tau^{-1}(e)$ of a given $e \in \tau(\Gamma')$ is the maximal subtree of $\Gamma'$ rooted in the unique minimal element of $\tau^{-1}(e)$. Thus $\tau$ is a morphism of trees compatible with the cochains $\sigma'$ and $\sigma$ in the sense (1.5), whence a morphism of $(A, x)$-labelled trees. $\square$

**Blow-downs of trees.** By definition, (see (1.5)), A morphism of trees $\tau : \Gamma' \to \Gamma$ factors through the retraction of a collection of maximal subtrees of $\Gamma'$, followed by an embedding. Therefore, for every element $e \in \Gamma' \setminus L(\Gamma')$ such that $\mathrm{Ch}_{\Gamma'}(e) \subset L(\Gamma')$, the image of the subtree $\Gamma'_e = (\uparrow e)_{\Gamma'}$ of $\Gamma'$ is either a subtree $\tau(\Gamma'_e)$ of $\Gamma$ isomorphic to $\Gamma'_e$ or the unique element $\tau(e) \in L(\Gamma)$. In the second case, $\tau$ factors through the morphism of trees

$$\tau_e : \Gamma' \longrightarrow \Gamma' \setminus \mathrm{Ch}_{\Gamma'}(e) \quad e' \mapsto \begin{cases} e' & \text{if } e' \in \Gamma' \setminus \mathrm{Ch}_{\Gamma'}(e) \\ e & \text{if } e' \in \mathrm{Ch}_{\Gamma'}(e) \end{cases}$$

**Definition 1.9.** Let $\gamma = (\Gamma, \sigma)$ be an $(A, x)$-labelled tree, and let $e \in \Gamma \setminus L(\Gamma)$ be an element such that $\mathrm{Ch}_{\Gamma}(e) \subset L(\Gamma)$. A *blow-down of the leaves at* $e$ is an $(A, x)$-labelled tree $\hat{\gamma}(e) = \left(\hat{\Gamma}(e), \hat{\sigma}(e)\right)$ with underlying tree $\hat{\Gamma}(e) = \Gamma \setminus \mathrm{Ch}_{\Gamma}(e)$ for which the morphism of trees $\tau_e$ above is a morphism of $(A, x)$-labelled trees $\tau_e : \gamma \to \hat{\gamma}(e)$. Since two labelled trees $\hat{\gamma}(e)$ with this property are isomorphic, the morphism $\tau_e$ itself will be usually referred to as *the* blow-down of the leaves at $e$.

As a consequence of the above discussion, we obtain the following description.

**Proposition 1.10.** *A morphism of $(A, x)$-labelled trees factors into a sequence of blow-downs of leaves followed by an embedding.*

**Equivalence of labelled trees.** An $(A, x)$-labelled tree $\gamma = (\Gamma, \sigma)$ rooted in $e_0$ is called *essential* if $\mathrm{Card}(\mathrm{Child}_{\Gamma}(e_0)) \neq 1$. The *essential subtree* $\mathrm{Es}(\Gamma)$ of a given tree $\Gamma$ is the maximal subtree of $\Gamma$ rooted either in the unique leaf of $\Gamma$ if $\Gamma$ is a chain, or in the first common ancestor $\tilde{e}_0$ of the leaves of $\Gamma$ otherwise. For instance, the tree $\Gamma$ of Figure 1 above is not essential, and its essential rooted subtree is the maximal subtree $(\uparrow e)_{\Gamma}$ of $\Gamma$ rooted in $e$. If an $(A, x)$-labelled tree $\gamma = (\Gamma, \sigma)$ is not essential then there exists $c \in A$ and an $\mathrm{Es}(\Gamma)$-compatible cochain $\mathrm{Es}(\sigma) = \{\tilde{\sigma}_i\}_{i=1,\ldots,n} \in A^n$ such that $\sigma_i = c + x^m \tilde{\sigma}_i$ for every $i = 1, \ldots, n$, where $m$ denotes the height of the root $\tilde{e}_0$ of the essential subtree $\mathrm{Es}(\Gamma)$ of $\Gamma$. A cochain $\mathrm{Es}(\sigma)$ with this property is called an *essential cochain* for $\gamma$, and we say that $\mathrm{Es}(\gamma) = (\mathrm{Es}(\Gamma), \mathrm{Es}(\sigma))$ is an *essential labelled subtree for* $\gamma$.

**Definition 1.11.** We say that two $(A,x)$-labelled trees $\gamma=(\Gamma,\sigma)$ and $\gamma'=(\Gamma',\sigma')$ are *equivalent* if there exist essential cochains $\mathrm{Es}\,(\sigma)$ and $\mathrm{Es}\,(\sigma')$ for $\gamma$ and $\gamma'$ respectively, an isomorphism of trees $\tau:\mathrm{Es}\,(\Gamma')\xrightarrow{\sim}\mathrm{Es}\,(\Gamma)$ and a pair $(a,b)\in A^{*}\times A$ such that

$$a\mathrm{Es}\,(\sigma')_{i}-\mathrm{Es}\,(\sigma)_{j(i)}+b\in x^{m_{j(i)}}A$$

whenever $e_{j(i),m_{j(i)}}\in L\,(\mathrm{Es}\,(\Gamma))$ is the image of $e'_{i,m'_{i}}\in L\,(\mathrm{Es}\,(\Gamma'))$ by $\tau$.

**Example 1.12.** By definition, two essential cochains for a given labelled tree $\gamma$ differ by the addition of a constant $b\in A$. Therefore, the essential labelled subtrees for $\gamma$ are all equivalent in the sense of definition (1.11).

## 2. Danielewski-Fieseler surfaces as fiber bundles

To fix the notation, we let $k$ be a field of caracteristic zero, and we let $(A,x)$ be a pair consisting either of a discrete valuation ring $A$ with uniformizing parameter $x$ and residue field $k$, or a polynomial ring $k\,[x]$ in one variable $x$. We let $X=\mathrm{Spec}\,(A)$, $x_{0}=\mathrm{div}\,(x)\in X$ , and we denote by $X_{*}=X\setminus\{x_{0}\}\simeq\mathrm{Spec}\,(A_{x})$ the open complement of $x_{0}$ in $X$.

**Definition 2.1.** A *Danielewski-Fieseler surface* (a DFS for short) with base $(X,x_{0})$ (or, equivalently, with base $(A,x)$) is an integral affine $X$-scheme $\pi:S\rightarrow X$ of finite type such that $S\mid_{X_{*}}$ is isomorphic to the trivial line bundle $\mathbb{A}^{1}_{X_{*}}$ over $X_{*}$ and such that the scheme-theoretic fiber $\pi^{-1}\,(x_{0})$ is nonempty and reduced, consisting of a disjoint union of curves isomorphic to the affine line $\mathbb{A}^{1}_{k}$. A DFS with base $(k\,[x]\,,x)$ is simply referred to as a *Danielewski surface*.

A *morphism of Danielewski-Fieseler surfaces* with base $(X,x_{0})$ is a birational $X$-morphism $\beta:S'\rightarrow S$ restricting to an $X_{*}$-isomorphism $\beta_{*}:S'\mid_{X_{*}}\xrightarrow{\sim}S\mid_{X_{*}}$.

Danielewski-Fieseler surfaces together with these morphisms form a sub-category $\mathfrak{D}_{(A,x)}$ (or $\mathfrak{D}_{(X,x_{0})}$) of the category $\left(\mathrm{Sch}_{/X}\right)$ of $X$-schemes.

**2.2.** The total space $S$ of DFS $\pi:S\rightarrow X$ is a smooth affine surface over $k$. Indeed, $X/k$ is itself affine and smooth, and the local criterion for flatness ([2, III,§5]) guarantees that $\pi$ is a faithfully flat morphism, whence a smooth morphism as its geometric fibers are regular. Note that in contrast with general DFS's, a Danielewski surface is an affine $k$-scheme of *finite type* since in this case $X\simeq\mathbb{A}^{1}_{k}$ by definition. In general, an arbitrary $\mathbb{A}^{1}$-fibration $\pi:S\rightarrow\mathbb{A}^{1}_{k}$ on a smooth affine surface $S$ does not give rise to a structure of Danielewski surface on $S$, but this does not prevent $S$ from being a Danielewski surface for another suitable $\mathbb{A}^{1}$-fibration. For instance, the Bandman and Makar-Limanov surface [1] $S\subset\mathrm{Spec}\,(k\,[x]\,[y,z,u])$ with equations

$$xz-y\,(y-1)=0,\quad yu-z^{2}=0,\quad xu-(y-1)\,z=0,$$

equipped with the $\mathbb{A}^{1}$-fibration $pr_{u}:S\rightarrow\mathrm{Spec}\,(k\,[u])$ is not a Danielewski surface with base $(k\,[u]\,,u)$. Indeed the fiber $pr_{u}^{-1}\,(0)$ is not reduced. However, it is a Danielewski surface $pr_{x}:S\rightarrow\mathrm{Spec}\,(k\,[x])$ with base $(k\,[x]\,,x)$.

Following Fieseler [7], DFS's with base $(X,x_{0})$ can be described as certain $\mathbb{A}^{1}$-bundles $\rho:S\rightarrow Y$ over a curve $\delta:Y\rightarrow X$ with an $n$-fold point over $x_{0}$. In this section, we give a new interpretation of Fieseler's 'cocycle construction'.

**Torsors under a line bundle.** Given an invertible sheaf $\mathcal{L}$ on a scheme $X$, the line bundle

$$\mathbf{V}(\mathcal{L}) = \mathbf{Spec}(\mathbf{S}(\mathcal{L})) = \mathbf{Spec}\left(\bigoplus_{n\geq 0}\mathcal{L}^n\right) \longrightarrow X$$

is equipped with the structure of a commutative group scheme for the group law $m : \mathbf{V}(\mathcal{L}) \times_X \mathbf{V}(\mathcal{L}) \to \mathbf{V}(\mathcal{L})$ defined by means of the $\mathcal{O}_X$-algebras homomorphism $\mathbf{S}(\mathcal{L}) \longrightarrow \mathbf{S}(\mathcal{L}) \otimes \mathbf{S}(\mathcal{L})$ induced by the diagonal homomorphism $\Delta : \mathcal{L} \to \mathcal{L} \oplus \mathcal{L}$. For instance, if $\mathcal{L} = \mathcal{O}_X$ then $\mathbf{V}(\mathcal{L})$ is simply the additive group scheme $\mathbb{G}_{a,X} = \mathbf{Spec}(\mathcal{O}_X[T])$.

**2.3.** A scheme $\pi : S \to X$ locally isomorphic to the trivial line bundle $\mathbb{A}^1_X$ is called an $\mathbb{A}^1$-*bundle*. If $S$ comes further equipped with an action of a line bundle $\mathbf{V}(\mathcal{L})/X$ for which there exists an open covering $\mathcal{U} = (X_i)_{i\in I}$ of $X$ such that $S\mid_{X_i}$ is equivariantly isomorphic to $\mathbf{V}(\mathcal{L})\mid_{X_i}$ acting on itself by translations for every $i \in I$, then we say that $S$ is a $\mathbf{V}(\mathcal{L})$-*torsor*. We recall [9, XI.4.7] that the set of isomorphism classes of $\mathbf{V}(\mathcal{L})$-torsors is a group isomorphic to $H^1(X, \mathcal{L}^\vee)$. Indeed, an automorphism of the symmetric $\mathcal{O}_X$-algebra $\mathbf{S}(\mathcal{L})$ is equivariant for the co-action of $\mathbf{S}(\mathcal{L})$ on itself by translations if and only if it is induced by an homomorphism of $\mathcal{O}_X$-modules $(g, \mathrm{Id}) : \mathcal{L} \to \mathcal{O}_X \oplus \mathcal{L}$, where $g \in \mathrm{Hom}_X(\mathcal{L}, \mathcal{O}_X)$. Therefore, given a $\mathbf{V}(\mathcal{L})$-torsor $\pi : S \to X$ which becomes trivial on $\mathcal{U}$, there exists a Čech cocycle $g = \{g_{ij}\}_{i,j\in I} \in C^1(\mathcal{U}, \mathcal{L}^\vee)$ such that $S$ is isomorphic to the scheme $\mathbf{W}(\mathcal{U}, \mathcal{L}, g)$ defined as the spectrum of the quasi-coherent $\mathcal{O}_X$-algebra obtained by gluing the symmetric algebras $\mathbf{S}(\mathcal{L}\mid_{X_i})$ over $X_{ij} = X_i \cap X_j$ by means of the $\mathcal{O}_{X_{ij}}$-algebras isomorphisms $\mathbf{S}(g_{ij}, \mathrm{Id})$ induced by the $\mathcal{O}_{X_{ij}}$-modules homomorphisms $(g_{ij}, \mathrm{Id}) \in \mathrm{Hom}_{X_{ij}}\left(\mathcal{L}\mid_{X_{ij}}, \mathcal{O}_{X_{ij}} \oplus \mathcal{L}\mid_{X_{ij}}\right)$, $i \neq j$. The isomorphism class of $\mathbf{W}(\mathcal{U}, \mathcal{L}, g)$ is simply the image in $H^1(X, \mathcal{L}^\vee) \simeq \check{H}^1(X, \mathcal{L}^\vee)$ of the class $[g] \in \check{H}^1(\mathcal{U}, \mathcal{L}^\vee)$ of $g \in C^1(\mathcal{U}, \mathcal{L}^\vee)$.

**2.4.** If the base scheme $X$ is integral, then every $\mathbb{A}^1$-bundle $\pi : S \to X$ admits a structure of a $\mathbf{V}(\mathcal{L})$-torsor for a suitable invertible sheaf $\mathcal{L}$ on $X$. Indeed, since $X$ is integral, the transition isomorphisms $\tau_{ij} = \tau_i \circ \tau_j^{-1}$ associated with a given collection of trivialisations $\tau_i : S\mid_{X_i} \xrightarrow{\sim} \mathrm{Spec}(\mathcal{O}_{X_i}[T])$, $i \in I$, are induced by automorphisms $T \mapsto \tilde{g}_{ij} + f_{ij}T$ of $\mathcal{O}_{X_{ij}}[T]$, where $\tilde{g}_{ij} \in \Gamma(X_{ij}, \mathcal{O}_X)$ and $f_{ij} \in \Gamma(X_{ij}, \mathcal{O}_X^*)$. For a triple of indices $i \neq j \neq k$, the identity $\tau_{ik} = \tau_{jk} \circ \tau_{ij}$ over $X_i \cap X_j \cap X_k$ guarantees that $(f_{ij})_{i,j\in I} \in C^1(\mathcal{U}, \mathcal{O}_X^*)$ is a Čech cocycle defining an invertible sheaf $\mathcal{L}$ on $X$, trivial on $\mathcal{U}$, with isomorphisms $\phi_i : \mathcal{L}\mid_{X_i} \xrightarrow{\sim} \mathcal{O}_{X_i}$, $i \in I$. By construction, $g = \left\{\tilde{g}_{ij} \cdot \phi_i\mid_{X_{ij}}\right\}_{i,j\in I} \in C^1(\mathcal{U}, \mathcal{L}^\vee)$ is a Čech cocycle for which $S$ is $X$-isomorphic to the $\mathbf{V}(\mathcal{L})$-torsor $\mathbf{W}(\mathcal{U}, \mathcal{L}, g)$.

**Schemes with an $n$-fold divisor.** Given a principal divisor $x_0 = \mathrm{div}(x)$ on an integral scheme $X$ and an integer $n \geq 1$, we let $\delta_n : X(n) = X(x_0, n) \to X$ be the scheme obtained by gluing $n$ copies $d_i : X_i(n) \xrightarrow{\sim} X$ of $X$ by the identity over the open subsets $X_i(n)_* = X_i(n) \setminus x_i(n)$, where $x_i(n) = d_i^{-1}(x_0)$. We denote by $\mathcal{U}_{(n)} = (X_i(n))_{i=1,\ldots,n}$ the canonical open covering of $X(n)$. For every $i \neq j$, we let $X_{ij}(n) = X_i(n) \cap X_j(n) \simeq X \setminus \{x_0\}$.

**Definition 2.5.** Given a multi-index $\mu = (\mu_1, \ldots, \mu_n) \in \mathbb{Z}^n$, we let

$$\mathcal{L}_{n,\mu} \quad = \quad \mathcal{O}_{X(n)}(\mu_1 x_1(n) + \ldots + \mu_n x_n(n))$$

be sub-$\mathcal{O}_{X(n)}$-module of the constant sheaf $\mathcal{K}_{X(n)}$ of rational functions on $X(n)$ generated by $(x \circ \delta_n)^{-\mu_i}$ on $X_i(n)$. The dual sheaf $\mathcal{L}_{n,\mu}^{\vee}$ of $\mathcal{L}_{n,\mu}$ is isomorphic to the sheaf $\mathcal{L}_{n,-\mu}$ corresponding to the multi-index $-\mu = (-\mu_1, \ldots, -\mu_n) \in \mathbb{Z}^n$. We denote by $s_\mu$ the canonical *rational* section of $\mathcal{L}_{n,\mu}$ corresponding to the constant section $1 \in \Gamma\left(X(n), \mathcal{K}_{X(n)}\right)$.

**Danielewski-Fieseler surfaces as torsors.** Given a DFS $\pi : S \to X$ with base $(X, x_0)$, we denote by $C_1, \ldots, C_n$ the connected component of the fiber $\pi^{-1}(x_0)$. We let $\delta_n : X(n) = X(x_0, n) \to X$ be as above, and we let $\rho_n : S \to X(n)$ be the unique $X$-morphism such that $\pi = \delta_n \circ \rho_n$, and such that $C_i = \rho_n^{-1}(x_i(n))$ for every $i = 1, \ldots, n$.

**Proposition 2.6.** *There exist a multi-index* $\mu = (\mu_1, \ldots, \mu_{n,}) \in \mathbb{Z}_{\geq 0}^n$ *such that* $\mu_i = 0$ *for at least one indice* $i$*, and a cocycle* $g \in C^1\left(\mathcal{U}_{(n)}, \mathcal{L}_{n,\mu}\right)$ *such that* $S$ *is* $X(n)$*-isomorphic to the* $\mathbf{V}(\mathcal{L}_{n,-\mu})$*-torsor* $\rho_n : \mathbf{W}\left(\mathcal{U}_{(n)}, \mathcal{L}_{n,-\mu}, g\right) \to X(n)$.

*Proof.* Since $X$ is either the affine line or the spectrum of a discrete valuation ring, the Picard group $\mathrm{Pic}(X)$ is trivial, and so every invertible sheaf on $X(n)$ is isomorphic to $\mathcal{L}_{n,-\mu}$ for a certain multi-index $\mu \in \mathbb{Z}_{\geq 0}^n$ such that $\mu_i = 0$ for at least one indice $i$. Moreover, every open subset $X_i(n) \simeq X$ of the covering $\mathcal{U}_{(n)}$ is affine. Therefore, a $\mathbf{V}(\mathcal{L}_{n,-\mu})$-torsor $\rho : W \to X(n)$ is isomorphic to $\mathbf{W}\left(\mathcal{U}_{(n)}, \mathcal{L}_{n,-\mu}, g\right)$ for a certain cocycle $g \in C^1(\mathcal{U}, \mathcal{L}_{n,\mu})$ representing its isomorphism class $c(W) \in H^1(X(n), \mathcal{L}_{n,\mu})$. So, by (2.4), it suffices to show that $\rho_n : S \to X(n)$ is an $\mathbb{A}^1$-bundle . This can be done in a similar way as in Lemma 1.2 in [7]. □

**2.7.** In view of the correspondence (2.4) between $\mathbb{A}^1$-bundles and $\mathbf{V}(\mathcal{L})$-torsors, a DFS $\mathbf{W}\left(\mathcal{U}_{(n)}, \mathcal{L}_{n,-\mu}, g\right)$ is $X$-isomorphic to the surface $S = \bigsqcup_{i=1}^n \mathbb{A}_{X_i}^1 / \sim$ obtained by gluing $n$ copies of $\mathbb{A}_{X_i}^1 = \mathrm{Spec}\,(A[T_i])$ of $\mathbb{A}_X^1$ by means of $A_x$-algebras isomorphisms $A_x[T_i] \xrightarrow{\sim} A_x[T_j]$, $T_i \mapsto \tilde{g}_{ij} + x^{\mu_i - \mu_j} T_j$, where $\tilde{g}_{ij} := x^{\mu_i} g_{ij} \in A_x$ for every $i \neq j$. In this way, we recover the "cocycle construction" of Fieseler [7] .

**2.8.** A general scheme $S = \mathbf{W}\left(\mathcal{U}_{(n)}, \mathcal{L}_{n,-\mu}, g\right)$ is not affine. For instance, if $n \geq 2$, then $\mathbb{A}_{X(n)}^1$ is not even separated. However, a similar argument as in Proposition 1.4 in [7] shows that $S$ is affine if and only if the corresponding *transition functions* $\tilde{g}_{ij} = x^{\mu_i} g_{ij}$ and $\tilde{g}_{ji} = -x^{\mu_j} g_{ij}$ above have a pole in $x_0$ for every $i \neq j$. This means equivalently that for every $i \neq j$, $g_{ij} \in \Gamma(X_{ij}(n), \mathcal{L}_{n,\mu})$ is not in the image of the differential

$$(\mathrm{res}_i - \mathrm{res}_j) : \Gamma(X_j(n), \mathcal{L}_{n,\mu}) \oplus \Gamma(X_i(n), \mathcal{L}_{n,\mu}) \to \Gamma(X_{ij}(n), \mathcal{L}_{n,\mu}).$$

In turn, this condition is satisfied if and only if for every $i \neq j$, $S\mid_{X_i(n) \cup X_j(n)}$ is a nontrivial $\mathbf{V}(\mathcal{L}_{n,-\mu})\mid_{X_i(n) \cup X_j(n)}$-torsor. This leads to the following criterion.

**Proposition 2.9.** *A scheme* $\mathbf{W}\left(\mathcal{U}_{(n)}, \mathcal{L}_{n,-\mu}, g\right)$ *is a DFS if and only if it restricts to a nontrivial* $\mathbf{V}(\mathcal{L}_{n,-\mu})\mid_U$*-torsor on every open subset* $U \simeq X(2)$ *of* $X$.

**Morphisms of Danielewski-Fieseler surfaces.** Given two DFS's $S' = \mathbf{W}\left(\mathcal{U}_{(n')}, \mathcal{L}_{n',-\mu'}, g'\right)$ and $S' = \mathbf{W}\left(\mathcal{U}_{(n')}, \mathcal{L}_{n',-\mu'}, g'\right)$ as in (2.6) and a morphism of DFS's $\beta : S' \to S$, we let $\alpha : X(n') \to X(n)$ be the unique $X$-morphism such that $\beta\left(\rho_{n'}^{-1}(x_i(n'))\right) \subset \rho_n^{-1}(\alpha(x_i(n)))$ for every $i = 1, \ldots, n'$. We denote by $\bar{g} \in C^1\left(\mathcal{U}_{(n')}, \alpha^* \mathcal{L}_{n,\mu}\right)$ the image of $\alpha^* g \in C^1\left(\alpha^{-1}\left(\mathcal{U}_{(n)}\right), \alpha^* \mathcal{L}_{n,\mu}\right)$ by the restriction maps between Čech complexes. We obtain a factorization $\beta = pr_S \circ \beta'$

$$\begin{tikzcd}
S' \arrow[r, "\beta'"] \arrow[d, "\rho_{n'}"'] & \tilde{S} = S \times_{X(n)} X(n') \simeq \mathbf{W}\left(\mathcal{U}_{(n')}, \alpha^* \mathcal{L}_{n,-\mu}, \bar{g}\right) \arrow[r, "pr_S"] \arrow[d, "p_2"'] & S \arrow[d, "\rho_n"'] \\
X(n') \arrow[r, equal] & X(n') \arrow[r, "\alpha"] & X(n)
\end{tikzcd}$$

where $\beta'$ is an $X(n')$-morphism restricting an isomorphism over $\delta_{n'}^{-1}(x_0)$. This is the case if and only if the homomorphisms $\mathbf{S}(\alpha^* \mathcal{L}_{n,-\mu}) \mid_{X_i(n')} \rightarrow \mathbf{S}(\mathcal{L}_{n',-\mu'}) \mid_{X_i(n')}$ corresponding to the restrictions $\beta'_i : S' \mid_{X_i(n')} \rightarrow \tilde{S} \mid_{X_i(n')}$ of $\beta'$ are induced by $\mathcal{O}_{X_i}$-modules homomorphisms $(c_i, \zeta_i) : \alpha^* \mathcal{L}_{n,-\mu} \mid_{X_i(n')} \rightarrow \mathcal{O}_{X_i(n')} \oplus \mathcal{L}_{n',-\mu'} \mid_{X_i(n')}$, $i = 1, \ldots, n'$, where $\zeta_i \in \mathrm{Hom}_{X_i(n')}(\alpha^* \mathcal{L}_{n,-\mu}, \mathcal{L}_{n',-\mu'})$ restricts to an isomorphism over $X_i(n') \backslash \{x_i(n')\}$, and where $c_i \in \Gamma(X_i(n'), \alpha^* \mathcal{L}_{n,\mu})$. Moreover, it follows from the definition of $S'$ and $\tilde{S}$ respectively that the $\beta_i$'s coincide on the overlaps $X_{ij}(n')$, $i \neq j$, if and only if the $\zeta_i$'s glue to a global section $\zeta \in \mathrm{Hom}_{X(n')}(\alpha^* \mathcal{L}_{n,-\mu}, \mathcal{L}_{n',-\mu'})$ such that $g' \circ \zeta = \bar{g} + \partial(c)$, where $\partial$ denotes the differential of the Čech complex $C^{\bullet}\left(\mathcal{U}_{(n')}, \alpha^* \mathcal{L}_{n,-\mu}\right)$. Summing up, we obtain the following characterization.

**Proposition 2.10.** *A morphism of DFS's $\beta : S' \rightarrow S$ exists if and only if there exists a data $(\alpha, \theta, c)$ consisting of:*
*1) an $X$-morphism $\alpha : X(n') \rightarrow X(n)$,*
*2) a section $\theta = {}^t\zeta \in Hom_{X(n')}(\mathcal{L}_{n',\mu'}, \alpha^* \mathcal{L}_{n,\mu})$ such that $Supp(\theta) \subset \delta_{n'}^{-1}(x_0)$,*
*3) a cochain $c \in C^0\left(\mathcal{U}_{(n')}, \alpha^* \mathcal{L}_{n,\mu}\right)$ such that $\theta(g') = \bar{g} + \partial(c)$.*
*Furthermore, $\beta$ is an isomorphism if and only if $\alpha$ and $\theta$ are.*

**Example 2.11.** Given a DFS $S = \mathbf{W}\left(\mathcal{U}_{(n)}, \mathcal{L}_{n,-\mu}, g\right)$, there exists an integer $h_0 \geq \max(\mu_i)$, depending on $g$ and $\mu$, such that for every $h \geq h_0$, $x^h s_{-\mu}$ is a regular section of $\mathcal{L}_{n,-\mu}$ defining an homomorphism of $\mathcal{O}_{X(n)}$-modules $\theta_{\mu,h} : \mathcal{L}_{n,\mu} \rightarrow \mathcal{O}_{X(n)}$ with the property that $\theta_{\mu,h}(g_{1j}) \in \Gamma\left(X_{1j}(n), \mathcal{O}_{X(n)}\right)$ extends to a section $\sigma_j$ of $\mathcal{O}_{X_j(n)}$ for every $j = 2, \ldots, n$. By (2.10), the data $(\delta_n, \theta_{\mu,h}, \sigma)$, where $\sigma = \{\sigma_1 = 0, \sigma_j\}_{j=2,\ldots,n} \in A^n$, corresponds to a morphism of DFS's $\psi : S \rightarrow \mathbb{A}^1_X$.

**Additive group actions on Danielewski-Fieseler surfaces.** As a torsor, a DFS $\pi : S = \mathbf{W}\left(\mathcal{U}_{(n)}, \mathcal{L}_{n,-\mu}, g\right) \xrightarrow{\rho} X(n) \overset{\delta_n}{\rightarrow} X$ comes equipped with an action $m_{n,\mu} : \mathbf{V}(\mathcal{L}_{n,-\mu}) \times_{X(n)} S \rightarrow S$ of the line bundle $\mathbf{V}(\mathcal{L}_{n,-\mu})$. Every nonzero section $s \in \Gamma(X(n), \mathcal{L}_{n,\mu})$ gives rise to a morphism of group schemes $\phi_s : \mathbb{G}_{a,X(n)} \rightarrow \mathbf{V}(\mathcal{L}_{n,-\mu})$, whence to a nontrivial action

$$m_s = m_{n,\mu} \circ (\phi_s \times \mathrm{Id}) : \mathbb{G}_{a,X(n)} \times_{X(n)} S \overset{\phi_s \times \mathrm{Id}}{\longrightarrow} \mathbf{V}(\mathcal{L}_{n,-\mu}) \times_{X(n)} S \overset{m_{n,\mu}}{\longrightarrow} S$$

of the additive group $\mathbb{G}_{a,X}$ by means of $X(n)$-automorphisms of $S$.

**Proposition 2.12.** *Every nontrivial $\mathbb{G}_{a,X}$-action on $S$ appear in this way.*

*Proof.* A connected component of $\pi^{-1}(x_0)$ is invariant under a $\mathbb{G}_{a,X}$-action. Therefore, a $\mathbb{G}_{a,X}$-action on $S$ lifts to a $\mathbb{G}_{a,X(n)}$-action $m : \mathbb{G}_{a,X(n)} \times_{X(n)} S \rightarrow S$. In turn, this action restricts on $S \mid_{X_i(n)} \simeq \mathbf{V}(\mathcal{L}_{n,-\mu}) \mid_{X_i(n)}$ to a $\mathbb{G}_{a,X_i(n)}$-action $m_i$. Thus there exists a nonzero section $s_i \in \Gamma(X_i(n), \mathcal{L}_{n,\mu}) \simeq \mathrm{Hom}_{X_i(n)}\left(\mathcal{L}_{n,-\mu}, \mathcal{O}_{X_i(n)}\right)$ such that the corresponding group co-action is induced by the $\mathcal{O}_{X_i(n)}$-modules homomorphism $(\mathrm{Id} \otimes 1 + s_i \otimes T) : \mathcal{L}_{n,-\mu} \mid_{X_i(n)} \rightarrow \mathbf{S}(\mathcal{L}_{n,-\mu}) \mid_{X_i(n)} \otimes_{\mathcal{O}_{X_i(n)}} \mathcal{O}_{X_i(n)}[T]$.

Clearly, these actions coincide on the overlaps $X_{ij}(n)$ if and only if the $s_i$'s glue to a section $s \in \Gamma(X, \mathcal{L}_{n,\mu})$ such that $m = m_s$. □

**Corollary 2.13.** *A DFS $\pi : S \to X$ admits a free $\mathbb{G}_{a,X}$-action if and only if its canonical sheaf is trivial.*

*Proof.* By construction, the canonical sheaf of $S = \mathbf{W}\left(\mathcal{U}_{(n)}, \mathcal{L}_{n,-\mu}, g\right)$ is isomorphic to $\rho_n^*\mathcal{L}_{n,-\mu}$. It is trivial if and only if $\mathcal{L}_{n,-\mu}$ is. On the other hand, the discussion above implies that $S$ admits a free $\mathbb{G}_{a,X}$-action if and only if $\mathcal{L}_{n,\mu} \simeq \mathcal{L}_{n,-\mu}^{\vee}$ is trivial. □

## 3. Danielewski-Fieseler surfaces and labelled rooted trees

Here we give a combinatorial description of DFS's by means of labelled trees. To state the main result of this section, we need the following definition.

**Definition 3.1.** *A relative Danielewski-Fieseler surface with base* $(A, x)$ is a morphism $\mathfrak{S} = \left(\psi : S \to \mathbb{A}_X^1\right)$ of DFS's with base $(A, x)$. A *morphism of relative DFS*'s is a morphism of DFS's $\beta : S' \to S$ such that $\psi' = \psi \circ \beta$.

This section is devoted to the proof of the following result.

**Theorem 3.2.** *The category $\overrightarrow{\mathfrak{D}}_{(A,x)}$ of relative DFS's with base $(A, x)$ is equivalent to the category $\mathcal{T}_{(A,x)}$ of $(A, x)$-labelled rooted trees.*

In the following subsections, we construct an equivalence of categories in the form of a covariant functor $\mathfrak{S} : \mathcal{T}_{(A,x)} \to \overrightarrow{\mathfrak{D}}_{/(A,x)}$.

**DFS's defined by labelled rooted trees.** Given an $(A, x)$-labelled tree $\gamma = (\Gamma, \sigma)$ with the leaves $e_{1,m_1}, \ldots, e_{n,m_n}$, we let $\mu = \mu(\gamma) = (\mu_i = h - m_i)_{i=1,\ldots,n} \in \mathbb{Z}_{\geq 0}^n$, where $h = h(\Gamma)$ denotes the height of $\Gamma$. Since $\mu_i \leq h$ for every $i = 1, \ldots, n$, the multiplication by the regular section $x^h s_{-\mu}$ of $\mathcal{L}_{n,-\mu}$ defines an homomorphism $\theta_{\mu,h} : \mathcal{L}_{n,\mu} = \mathcal{L}_{n,-\mu}^{\vee} \to \mathcal{O}_{X(n)}$ restricting to an isomorphism over $\delta_n^{-1}(X_*)$. We let $g(\gamma) \in C^1\left(\mathcal{U}_{(n)}, \mathcal{L}_{n,\mu}\right)$ be the unique cocycle such that $\theta_{\mu,h}(g(\gamma)) = \partial(\sigma) \in C^1\left(\mathcal{U}_{(n)}, \mathcal{O}_{X(n)}\right)$. If $n = 1$, then the scheme $\pi_\gamma : S(\gamma) = \mathbf{W}\left(\mathcal{U}_{(n)}, \mathcal{L}_{n,-\mu}, g(\gamma)\right) \to X$ corresponding to this data is isomorphic to $\mathbb{A}_X^1$. Otherwise, if $n \geq 2$, then, by (1.5), the transition functions $\tilde{g}_{ij} = x^{\mu_i} g_{ij} = x^{-m_i}(\sigma_j - \sigma_i) \in A_x$ and $\tilde{g}_{ji}$ have a pole at $x_0$. Thus $S(\gamma)$ is a DFS by virtue of (2.8). The morphism of DFS's $\mathfrak{S}(\gamma) = \left(\psi_\gamma : S(\gamma) \to \mathbb{A}_X^1\right)$ defined by the data $(\delta_n, \theta_{\mu,h}, \sigma)$ is called the *canonical morphism associated with* $\gamma$. The following result completes the first part of the proof of theorem (3.2).

**Proposition 3.3.** *Every relative DFS is isomorphic to $\mathfrak{S}(\gamma)$ for a suitable tree $\gamma$.*

*Proof.* A morphism of DFS's $\mathfrak{S} = \left(\psi : S = \mathbf{W}\left(\mathcal{U}_{(n)}, \mathcal{L}_{n,-\mu}, g\right) \to \mathbb{A}_X^1\right)$ is given by a data $(\delta_n, \theta = a\theta_{\mu,h}, \sigma)$, where $a \in A^*$ and $h \geq \max(\mu_i)$, such that $\theta(g) = \partial(\sigma)$. If $n \geq 2$ then $m_i = h - \mu_i \geq 1$ for every $i = 1, \ldots, n$. Indeed, otherwise there exists an indice $i$ for which $\theta$ induces an isomorphism $\theta_i : \mathcal{L}_{n,\mu} \mid_{X_i(n)} \xrightarrow{\sim} \mathcal{O}_{X_i(n)}$. Thus, for every $j \neq i$, $g_{ij} \in \Gamma(X_{ij}(n), \mathcal{L}_{n,\mu})$ extends to a section of $\mathcal{L}_{n,\mu} \mid_{X_i(n)}$ as $\theta(g_{ij}) = \sigma_j \mid_{X_i(n)} - \sigma_i \mid_{X_i(n)} \in \Gamma\left(X_{ij}(n), \mathcal{O}_{X(n)}\right)$ extends to a section of $\mathcal{O}_{X_i(n)}$. This implies that $S \mid_{X_i(n) \cup X_j(n)}$ is a trivial torsor, in contradiction with (2.9). For the same reason, $d_{ij} = \operatorname{ord}_{x_0}(\sigma_j - \sigma_i) < \min(m_i, m_j)$ for every $i \neq j$. Since $\min(d_{ij}, d_{ik}) = \min(d_{ij}, d_{jk})$ for every triple of indices $i, j$ and $k$, we deduce from

(1.7) that the data $\left(n, (m_i)_{i=1,\ldots,n}, (d_{ij})_{i,j=1,\ldots,n}, \sigma\right)$ corresponds to an $(A,x)$-labelled tree $\gamma = (\Gamma, \sigma)$. Finally, the data $\left(\mathrm{Id}_{X(n)}, a\mathrm{Id}_{\mathcal{L}_{n,-\mu}}, 0\right)$ defines an isomorphism of DFS's $\phi : S \xrightarrow{\sim} S(\gamma)$ such that $\psi_\gamma = \psi \circ \phi^{-1} : S(\gamma) \to \mathbb{A}^1_X$. □

**Corollary 3.4.** *Every DFS is $X$-isomorphic to $S(\gamma)$ for a suitable tree $\gamma$.*

*Proof.* Indeed, by (2.11), every DFS $S$ admits a morphism of DFS's $\mathfrak{S} = \left(\psi : S \to \mathbb{A}^1_X\right)$. □

**Example 3.5.** Given three integers $r \geq 0$, $m \geq 1$ and $n \geq 1$, we consider an $(A,x)$-labelled tree $\gamma = (\Gamma_{r,m,n}, \sigma)$ with the following underlying tree

$\Gamma_{r,m,n} =$ 

Since there exists $b \in A$ and a cochain $\tilde{\sigma} \in A^n$ such that $\sigma_i = b + x^r \tilde{\sigma}_i$ for every $i = 1, \ldots, n$, the corresponding DFS $\pi_\gamma : S(\gamma) \to X$ is obtained by gluing $n$ copies $\mathrm{Spec}(A[T_i])$ of $\mathbb{A}^1_X$ by means of the $A_x$-algebras isomorphisms

$$A_x[T_i] \xrightarrow{\sim} A_x[T_j], \quad T_i \mapsto x^{-m}(\tilde{\sigma}_j - \tilde{\sigma}_i) + T_i, \quad i \neq j.$$

Since $\tilde{\sigma}_j - \tilde{\sigma}_i \in A \setminus xA$ for every $i \neq j$, the local sections $\tilde{\sigma}_i + x^m T_i \in A[T_i]$, $i = 1, \ldots, n$, glue to a global one $s_1 \in B = \Gamma\left(X, \pi_{\gamma *}\mathcal{O}_{S(\gamma)}\right)$ which distinguishes the irreducible components of $\pi_\gamma^{-1}(x_0)$. Letting $P = \prod_{i=1}^n (y - \tilde{\sigma}_i) \in A[y]$, the rational section $x^{-m}P(s_1) \in B \otimes_A A_x$ extends to a regular section $s_2 \in B$, inducing a coordinate function on every irreducible component of $\pi_\gamma^{-1}(x_0)$. By construction, the $A$-algebras homomorphism $A[y,z] \to B$, $y \mapsto s_1$, $z \mapsto s_2$ defines a closed embedding $\phi : S(\gamma) \hookrightarrow \mathbb{A}^2_X = \mathrm{Spec}(A[y,z])$ which induces an $X$-isomorphism between $S(\gamma)$ and the surface

$$\pi : S_{P,m} = \mathrm{Spec}\left(A[y,z] / (x^m z - P(y))\right) \longrightarrow X.$$

In this coordinates, the canonical morphism $\psi_\gamma : S(\gamma) \to \mathbb{A}^1_X$ is given as the restriction on $S_{P,m}$ of the $X$-morphism $\mathbb{A}^2_X \to \mathbb{A}^1_X$, $(y,z) \mapsto x^r y + b$.

**Morphism of DFS's defined by a morphism labelled rooted trees.** Given two $(A,x)$-labelled trees $\gamma' = (\Gamma', \sigma')$ and $\gamma = (\Gamma, \sigma)$, we equip the corresponding DFS's $S(\gamma') = \mathbf{W}\left(\mathcal{U}_{(n')}, \mathcal{L}_{n',-\mu'}, g'\right)$ and $S(\gamma) = \mathbf{W}\left(\mathcal{U}_{(n)}, \mathcal{L}_{n,-\mu}, g\right)$ with their canonical morphisms $\psi_{\gamma'}$ and $\psi_\gamma$ respectively. By (1.4), the image of a leaf $e'_{i,m'_i}$ of $\Gamma'$ by a morphism of $(A,x)$-labelled tree $\tau : \gamma' \to \gamma$ is a leaf $e_{j(i),m_{(j)}}$ of $\Gamma$ such that $m'_i \geq m_{j(i)}$, and $\sigma'_i - \sigma_{j(i)} \in x^{m_{j(i)}}A$ for every $i = 1, \ldots, n'$. Letting $\alpha : X(n') \to X(n)$ be the unique $X$-morphism such that $\alpha(x_i(n')) = x_{j(i)}(n)$ for every $i = 1, \ldots, n$, we conclude that the invertible sheaf $\alpha^*\mathcal{L}_{n,-\mu}$ is isomorphic to $\mathcal{L}_{n',-\nu}$, where $\nu = \left(\mu_{j(i)}\right)_{i=1,\ldots,n'} \in \mathbb{Z}^{n'}_{\geq 0}$. The multiplication by the regular section $x^{h(\Gamma')-h(\Gamma)} s_{\nu-\mu'} \in \Gamma\left(X(n'), \mathcal{L}_{n',\nu-\mu'}\right)$ defines an homomorphism of $\mathcal{O}_{X(n')}$-modules $\theta : \mathcal{L}_{n',\mu'} \to \alpha^*\mathcal{L}_{n,\mu} \simeq \mathcal{L}_{n',\nu}$ such that $\theta_{\mu',h'} = \alpha^*\theta_{\mu,h} \circ \theta$. By construction, there exists a unique cochain $\sigma'' \in C^0\left(\mathcal{U}_{(n')}, \mathcal{L}_{n',\nu}\right)$ such that

$\alpha^*\theta_{\mu,h}(\sigma'') = \{\sigma'_i - \sigma_{j(i)}\}_{i=1,\ldots n'} \in C^0\left(\mathcal{U}_{(n')}, \mathcal{O}_{X(n')}\right)$. Since $\alpha^*\theta_{\mu,h}$ restricts to an isomorphism over $\delta_{n'}^{-1}(X_*)$, we conclude that $\theta(g_{\sigma'}(\gamma')) = \overline{g_\sigma(\gamma)} + \partial(\sigma'')$, where $\overline{g_\sigma(\gamma)} \in C^1\left(\mathcal{U}_{(n')}, \mathcal{L}_{n',\nu}\right)$ denotes the image of $\alpha^* g_\sigma(\gamma) \in C^1\left(\alpha^{-1}\mathcal{U}_{(n)}, \alpha^*\mathcal{L}_{n,\mu}\right)$ by the restriction maps between Čech complexes. By (2.10), the data $(\alpha, \theta, \sigma'')$ defines a morphism of DFS's $\beta_\tau : S(\gamma') \to S(\gamma)$ such that $\psi_{\gamma'} = \psi_\gamma \circ \beta_\tau$, *i.e.* a morphism $\beta_\tau : \mathfrak{S}(\gamma') \to \mathfrak{S}(\gamma)$ in $\overrightarrow{\mathfrak{D}}_{(A,x)}$. We say that $\beta_\tau$ is the morphism of relative DFS's defined by $\tau : \gamma' \to \gamma$.

**Example 3.6.** We consider the blow-down $\tau_e : \gamma' \to \gamma$ of the leaves at $e$ between the following $(k[x], x)$-labelled trees $\gamma' = (\Gamma', \sigma')$ and $\gamma = (\Gamma, \sigma)$.



We let $s'_1 \in B' = \Gamma\left(X, (\pi_{\gamma'})_* \mathcal{O}_{S(\gamma')}\right)$ and $s_1 \in B = \Gamma\left(X, (\pi_\gamma)_* \mathcal{O}_{S(\gamma)}\right)$ be the sections corresponding to the canonical morphisms $\psi_{\gamma'} : S(\gamma') \to \mathbb{A}^1_X = \operatorname{Spec}(k[x][y])$ and $\psi_\gamma : S(\gamma) \to \mathbb{A}^1_X$ respectively. By (3.5), $S(\gamma)$ is isomorphic to the surface $S = \left\{xz - y\left(y^2 - 1\right) = 0\right\} \subset \mathbb{A}^2_X = \operatorname{Spec}(k[x][y,z])$ via the embedding induced by the $k[x]$-algebra homomorphism $k[x][y,z] \mapsto B$, $(y,z) \mapsto \left(s_1, s_2 = x^{-1}s_1\left(s_1^2 - 1\right)\right)$. A similar argument shows that $S(\gamma')$ is isomorphic to the Bandman and Makar-Limanov surface $S' \subset \mathbb{A}^3_X = \operatorname{Spec}(k[x][y,z,u])$ with equations

$$xz - y\left(y^2 - 1\right) = 0, \quad yu - z\left(z^2 - 1\right) = 0, \quad xu - \left(y^2 - 1\right)\left(z^2 - 1\right) = 0,$$

via the embedding induced by the $k[x]$-algebras homomorphism $k[x][y,z,u] \to B$

$$(y,z,u) \mapsto \left(s'_1, s'_2 = x^{-1}s'_1\left(\left(s'_1\right)^2 - 1\right), s'_3 = x^{-1}\left(\left(s'_1\right)^2 - 1\right)\left(\left(s'_2\right)^2 - 1\right)\right)$$

In these coordinates, the morphism $\beta_{\tau_e} : S(\gamma') \to S(\gamma)$ defined by the blow-down $\tau_e$ coincide with the restriction of the projection $\mathbb{A}^3_X \to \mathbb{A}^2_X$, $(x,y,z,u) \mapsto (x,y,z)$.

**3.7.** The correspondence $\gamma \mapsto \mathfrak{S}(\gamma)$, $(\tau : \gamma' \to \gamma) \mapsto \mathfrak{S}(\tau) = (\beta_\tau : \mathfrak{S}(\gamma') \to \mathfrak{S}(\gamma))$ defines a covariant functor $\mathfrak{S} : \mathcal{T}_{(A,x)} \to \overrightarrow{\mathfrak{D}}_{(A,x)}$. It follows from (3.3) that $\mathfrak{S}$ is essentially surjective. The following result shows that $\mathfrak{S}$ is fully faithful, whence completes the proof of theorem (3.2).

**Proposition 3.8.** *Every morphism $\beta : \mathfrak{S}(\gamma') \to \mathfrak{S}(\gamma)$ coincides with a unique morphism $\beta_\tau$ defined by a morphism of $(A,x)$-labelled trees $\tau : \gamma' \to \gamma$.*

*Proof.* By (2.10), $\beta$ is determined by a data $(\alpha, \theta, \sigma'')$ such that $\alpha(x_i(n')) = x_{j(i)}(n)$, $i = 1, \ldots, n'$, $\theta_{\mu',h} = \alpha^*\theta_{\mu,h} \circ \theta$, $\sigma' = \alpha^*\sigma + \alpha^*\theta_{\mu,h}(\sigma'')$, and such that

$$\theta(g_{\sigma'}(\gamma')) = \overline{\alpha^* g_\sigma(\gamma)} + \partial(\sigma'') \in C^1\left(\mathcal{U}_{(n')}, \alpha^*\mathcal{L}_{n,\mu}\right).$$

Since $\alpha^*\mathcal{L}_{n,\mu}$ is isomorphic to $\mathcal{L}_{n',\nu}$, where $\nu = \left(\mu_{j(i)}\right)_i \in \mathbb{Z}^{n'}_{\geq 0}$, we conclude that $\theta$ is the multiplication by $x^{h'-h}s_{\nu-\mu'} \in \Gamma(X(n'), \mathcal{L}_{n',\nu-\mu'})$. Thus $m'_i \geq m_{j(i)}$ and

$\tilde{\sigma}''_i = \alpha^* \theta_{\mu,h}\left(\sigma''_i\right) \in x^{h-\mu_{j(i)}} A = x^{m_{j(i)}} A$ for every $i = 1, \ldots, n'$. Therefore, the formulas

$$\Gamma'_i \ni e'_{i,k} \mapsto \tau_i\left(e'_{i,k}\right) = e_{j(i),\min\left(k,m_{j(i)}\right)} \in \Gamma_{j(i)}, \quad i = 1, \ldots, n' \; k = 0, \ldots, m'_i,$$

(see (1) for the notation) define a collection of morphisms of trees $\tau_i : (\Gamma'_i, \sigma'_i) \to \left(\Gamma_{j(i)}, \sigma_{j(i)}\right)$ in $\mathcal{T}_{(A,x)}$. Moreover, we deduce that

$$\text{(2)} \qquad \operatorname{ord}_{x_0}\left(\sigma'_{i'} - \sigma'_i\right) = \begin{cases} \operatorname{ord}_{x_0}\left(\tilde{\sigma}''_{i'} - \tilde{\sigma''}_i\right) \geq m_{j(i)} & \text{if } j(i) = j(i') \\ \operatorname{ord}_{x_0}\left(\sigma_{j(i')} - \sigma_{j(i)}\right) & \text{otherwise} \end{cases} .$$

Since $\sigma'$ and $\sigma$ are compatible with $\Gamma'$ and $\Gamma$ respectively, (2) guarantees that the conditions of (1.8) are satisfied. Thus there exists a unique morphism $\tau : \gamma' \to \gamma$ in $\mathcal{T}_{(A,x)}$ such that $\tau \mid_{\gamma'_i} = \tau_i$ for every $i = 1, \ldots, n'$. By construction, $\beta = \beta_\tau$. □

**Corollary 3.9.** *For every morphism of DFS's $\beta : S' \to S$ there exist a morphism of $(A,x)$-labelled trees $\tau : \gamma' \to \gamma$ and a commutative diagram*

$$\begin{tikzcd} S' \arrow[r, "\beta"] \arrow[d, "\phi'"', "\wr"] & S \arrow[d, "\phi", "\wr"'] \\ S(\gamma') \arrow[r, "\beta_\tau"] & S(\gamma) \end{tikzcd}$$

*Proof.* Once a morphism of DFS's $\psi : S \to \mathbb{A}^1_X$ is chosen (see (2.11)), $\left(\psi : S \to \mathbb{A}^1_X\right)$ and $\left(\psi' = \psi \circ \beta : S' \to \mathbb{A}^1_X\right)$ are objects of $\overrightarrow{\mathfrak{D}}_{(A,x)}$, whereas $\beta : S' \to S$ corresponds to morphism in $\overrightarrow{\mathfrak{D}}_{(A,x)}$. So the result follows from (3.3) and (3.8). □

**Reading isomorphism classes of Danielewski-Fieseler surfaces from trees.** To decide when two DFS's are $X$-isomorphic, we have the following criterion.

**Theorem 3.10.** *Two $(A,x)$-labelled trees define $X$-isomorphic DFS's if and only if they are equivalent (see (1.11) for the definition).*

*Proof.* We first observe that if $\gamma' = (\Gamma', \sigma')$ is an essential labelled subtree for $\gamma = (\Gamma, \sigma)$ then the DFS's $S(\gamma)$ and $S(\gamma')$ are $X$-isomorphic. Indeed, there exists $m \in \mathbb{Z}_{\geq 0}$ and $b \in A$ such that $m_i = m'_i + m$ and $\sigma_i = b + x^m \sigma'_i$ for every $i = 1, \ldots, n$. Thus $\mu(\gamma') = \mu(\gamma) = \mu \in \mathbb{Z}^n_{\geq 0}$, $g(\gamma) = g(\gamma') \in C^1\left(\mathcal{U}_{(n)}, \mathcal{L}_{n,\mu}\right)$, and so, $S(\gamma) \simeq S(\gamma')$. Therefore, it suffices to prove the assertion for DFS's defined by essential labelled trees $\gamma = (\Gamma, \sigma)$ and $\gamma' = (\Gamma', \sigma')$. If $\gamma$ and $\gamma'$ are equivalent then $h = h(\Gamma) = h(\Gamma')$ and there exists a permutation $j$ of $\{1, \ldots, n\}$ such that $\mu'_i = \mu_{j(i)}$ for every $i = 1, \ldots, n$. Moreover, there exist a pair $(a, b) \in A^* \times A$ and a cochain $\tilde{\sigma} = \{\tilde{\sigma}_i\}_{i=1,\ldots,n} \in A^n$ such that $\sigma'_i = a\sigma_{j(i)} + b + x^{m_{j(i)}} \tilde{\sigma}_{j(i)}$ for every $i = 1, \ldots, n$. Letting $\sigma'' = \{\sigma''_i = \sigma'_i - b\}_{i=1,\ldots,n}$, the same argument as above shows that $S(\gamma')$ is $X$-isomorphic to the DFS defined by the labelled tree $\gamma'' = (\Gamma', \sigma'')$. Thus, by replacing $S(\gamma')$ by $S(\gamma'')$ if necessary, we can suppose from now on that $b = 0$. We let $\alpha \in \operatorname{Aut}_X(X(n))$ be the unique $X$-automorphism such that $\alpha(x_i(n)) = x_{j(i)}(n)$, and we let $c \in C^0\left(\mathcal{U}_{(n)}, \alpha^* \mathcal{L}_{n,\mu}\right)$ be the unique cochain such that $\theta_{\mu',h}(c) = \left\{a^{-1} x^{m_{j(i)}} \tilde{\sigma}_{j(i)}\right\}_{i=1,\ldots,n}$. Since $\mathcal{L}_{n,\mu'} \simeq \alpha^* \mathcal{L}_{n,\mu}$ and $a^{-1} g(\gamma') = \alpha^* g(\gamma) + \partial(c)$, we conclude that the data $\left(\alpha, a\mathrm{Id}_{\mathcal{L}_{n,\mu'}}, c\right)$ defines an isomorphism of DFS's $\beta : S(\gamma') \xrightarrow{\sim} S(\gamma)$. Conversely, an $X$-isomorphism $\beta : S' =$

$S(\gamma') \overset{\sim}{\to} S = S(\gamma)$ exists if and only if the trees $\gamma$ and $\gamma'$ have the same number $n$ of leaves, and there exists a data

$$(\alpha, \theta, c) \in \mathrm{Aut}_X(X(n)) \times \mathrm{Isom}_{X(n)}(\mathcal{L}_{n,\mu'}, \alpha^*\mathcal{L}_{n,\mu}) \times C^0\left(\mathcal{U}_{(n)}, \alpha^*\mathcal{L}_{n,\mu}\right)$$

such that $\theta(g(\gamma')) = \alpha^* g(\gamma) + \partial(c)$. By replacing $S$ by the $X$-isomorphic DFS $\alpha^* S$, we can suppose that $\alpha = \mathrm{Id}_{X(n)}$. We can also suppose that $h' = h(\Gamma') \geq h = h(\Gamma)$. Then $\mathcal{L}_{n,\mu'}$ and $\mathcal{L}_{n,\mu}$ are isomorphic if and only if $m_i' = h' - \mu_i' = h - \mu_i + m = m_i + m$ for some $m \in \mathbb{Z}_{\geq 0}$. The identity $\theta(g(\gamma')) = g(\gamma) + \partial(c)$ is satisfied if and only there exist a pair $(a, b) \in A^* \times A$ such that $\tilde{\sigma}_i := \sigma_i + \theta_{\mu,h}(c_i) = a x^m \sigma_i' + b \in A$ for every $i = 1, \ldots, n$. By definition, $\theta_{\mu,h}(c_i) \in x^{m_i} A \subset xA$ for every $i = 1, \ldots, n$. Moreover, since $\Gamma$ is essential, there exist $i \neq j$ such that $\sigma_j - \sigma_i \in A \setminus xA$. Thus $m = 0$, and so $\mathrm{ord}_{x_0}(\sigma_j - \sigma_i) = \mathrm{ord}_{x_0}\left(\sigma_j' - \sigma_i'\right)$ for every $i \neq j$. Letting $\tilde{\sigma} = \{\tilde{\sigma}_i\}_{i=1,\ldots,n}$, we conclude that the trees $\gamma'$ and $\tilde{\gamma} = (\Gamma, \tilde{\sigma})$ are equivalent. This completes the proof as $\gamma$ and $\tilde{\gamma}$ are isomorphic, whence equivalent. □

**Example 3.11.** The following $(k[x], x)$-labelled trees $\gamma_t = (\Gamma, \sigma_t)$, where $t \in k$,

$\gamma_t =$



define a family of DFS's $\pi_t : S_t = S(\gamma_t) \to X = \mathrm{Spec}(k[x])$. A similar argument as in (3.5) shows that $S_t$ is $X$-isomorphic to the fiber $pr_v^{-1}(t)$ of the threefold $pr_v : \mathcal{X} \subset \mathrm{Spec}(k[x][y, z, u, v]) \to \mathrm{Spec}(k[v])$ with equations

$$xz - y(y-1) = 0, \quad yu - z(z-v) = 0, \quad xu - y(z-v) = 0.$$

These DFS's $S_t$ are two by two non $X$-isomorphic as the trees $\gamma_t$ are two by two nonequivalent. However, there are all isomorphic as abstract schemes. Indeed, a similar argument as in (4.6) below shows that there exists a projective surface $\bar{S}_t$, obtained from the Hirzebruch surface $\bar{p}_1 : \mathbb{F}_1 = \mathbb{P}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(-1)) \to \mathbb{P}^1_k$ by first blowing two distinct points $y_1, y_2 \in F_0 = \bar{p}_1^{-1}(0)$ with exceptional divisors $E$ and $F$, and then blowing-up a point $y_3 \in E \setminus F_0'$, and an open embedding of $S_t$ in $\bar{S}_t$ as the complement of the union of the strict transforms of $E$, $F_0$, $F_\infty = \bar{p}_1^{-1}(\infty)$ and of a section $C$ of $\bar{p}_1$ 'at infinity'. By successively blowing-down the strict transforms of $C$, $F_0$ and $E$, we realize $S_t$ as the complement in $\mathbb{F}_1$ of a section $D_t$ of $\bar{p}_1$ with self-intersection $\left(D_t^2\right) = 3$. By a result of Gizatullin-Danilov [8], the isomorphism class of $\mathbb{F}_1 \setminus D$ as an abstract scheme does not depend on the choice of a section $D$ such that $\left(D^2\right) = 3$. Therefore, the surfaces $S_t$, $t \in k$, are all isomorphic. Since they are not $X$-isomorphic, we deduce that a surface $S \simeq \mathbb{F}_1 \setminus D$ as above comes equipped with a family $q_t : S \to X_t$ of structures of DFS over different bases $X_t \simeq \mathbb{A}^1_k$, $t \in k$, such that the general fibers of $q_t$ and $q_{t'}$ do not coincide whenever $t \neq t'$.

## 4. DANIELEWSKI-FIESELER SURFACES AND AFFINE MODIFICATIONS

Here we exploit the structure of rooted trees to give another description of morphisms of DFS's. As a consequence, we obtain a canonical procedure to construct an open embedding of a DFS $\pi : S \to X$ into a projective $X$-scheme $\bar{\pi} : \bar{S} \to X$.

**Morphisms of Danielewski-Fieseler surfaces as affine modifications.** In what follows, we freely use the results about affine modifications, referring the reader to [11] for complete proofs. However, we recall the following definition.

**Definition 4.1.** Suppose we are given an affine scheme $V = \operatorname{Spec}(B)$, a nonzero divisor $f \in B$ an an ideal $I \subset B$ containing $f$. The *proper transform* $D_f^{pr}$ *of* $\operatorname{div}(f)$ in $\bar{V} = \operatorname{Proj}(B[It])$ is the set of prime ideals $\mathfrak{p} \in \bar{V}$ such that $ft \in \mathfrak{p}$. The affine scheme $V' = \bar{V} \setminus \operatorname{Supp}\left(D_f^{pr}\right) \simeq \operatorname{Spec}(B[It]/(1-ft))$ is called the *affine modification of $V$ with the locus $(I, f)$.*

**Definition 4.2.** A morphism of DFS's $\beta : S' \to S$ is called a *fibered modification* if there exist an $(A, x)$-labelled tree $\gamma' = (\Gamma', \sigma')$, a blow-down $\tau_{e'} : \gamma' \to \gamma$ of the leaves at a certain $e' \in \Gamma'$ (see (1.9) for the definition) and a commutative diagram

$$\begin{tikzcd}
S' \arrow[r, "\beta"] \arrow[d, "\wr"'] & S \arrow[d, "\wr"] \\
S(\gamma') \arrow[r, "\beta_{\tau'_e}"] & S(\gamma)
\end{tikzcd}$$

**4.3.** Consider a blow-down $\tau_{e'} : \gamma' \to \gamma$ of the leaves $e'_1, \ldots, e'_r$ at $e' \in \Gamma'$. For every $e'_i \in \operatorname{Leaves}(\Gamma) \setminus \{e'_1, \ldots, e'_r\}$, $e_{j(i)} = \tau_{e'}(e'_i)$ is a leaf of $\gamma'$, and $\tau_{e'}$ restricts to an isomorphism of labelled chains $(\downarrow e'_i)_{\gamma'} \xrightarrow{\sim} \left(\downarrow e_{j(i)}\right)_{\gamma}$. Therefore, the morphism $\beta_{\tau_{e'}} : S' = S(\gamma') \to S = S(\gamma)$ induces isomorphisms $S' \mid_{X_i(n')} \xrightarrow{\sim} S \mid_{X_{j(i)}(n)}$ between the corresponding open subsets of $S'$ and $S$ respectively. On the other hand, the image of the labelled subtree $\gamma(e') = (\downarrow e')_{\gamma'} \cup \operatorname{Ch}_{\Gamma'}(e')$ of $\gamma'$ by $\tau_{e'}$ is the subtree $\left(\downarrow_{\tau_{e'}(e')}\right)_{\gamma}$ of $\gamma$. By (3.5), there exists an $X$-isomorphism between $\tilde{S} = S' \mid_{X_1(n') \cup \cdots \cup X_r(n')}$ and a DFS $S_P = \operatorname{Spec}(A[y,z]/(xz - P(y)))$ for a certain polynomial $P \in A[y]$ of degree $r$ whose residue class $\bar{P} \in A[y]/xA[y] \simeq k[y]$ has $r$ simple roots. In this coordinates, the restriction $\beta : \tilde{S} \to S \mid_{X_{j(i)}(n)} \simeq \mathbb{A}^1_X$ coincides with the first projection $p_y : S_P \to \mathbb{A}^1_X = \operatorname{Spec}(A[y])$. Therefore, $\tilde{S}$ is isomorphic to the affine modification of $\mathbb{A}^1_X$ with locus $(I, f) = ((x, P(y)), x)$. Since $(x, P(y))$ is a regular sequence in $A[y]$, the Rees algebra $A[y][It]$ is isomorphic to $A[y][u,v]/(xv - P(y)u)$ via the map $u \mapsto xt$, $v \mapsto P(y)t$. Therefore, the proper transform $D_x^{pr} = \{xt = u = 0\}$ of the line $L = \operatorname{div}(x) \subset \mathbb{A}^1_X$ by the blow-up morphism $\bar{\beta} : \bar{S}_P = \operatorname{Proj}_A(A[y][It]) \to \mathbb{A}^1_X$ coincides with the usual strict transform $L'$ of $L$. This leads to the following description.

**Proposition 4.4.** *Given a fibered modification $\beta : S' \to S$, there exists a unique irreducible component $C$ of $\pi^{-1}(x_0)$ such that $\beta\left(\beta^{-1}(C)\right)$ is supported at a 0-dimensional closed subscheme $Y \subset C$. Letting $\bar{\beta} : \bar{S}' \to S$ be the blow-up of $Y$ equipped with its reduced structure, $S'$ is isomorphic to the complement in $\bar{S}'$ of the strict transform of $C$, and the irreducible components of $(\pi')^{-1}(x_0)$ contained in $\beta^{-1}(C)$ coincides with the intersections of the exceptional divisors of $\bar{\beta}$ with $S'$.*

As a consequence of (1.10) and (3.2), we obtain the following result.

**Theorem 4.5.** *Every morphism of Danielewski-Fieseler surfaces factors into a finite sequence of fibered modifications followed by an open embedding.*

**Local completions of Danielewski-Fieseler surfaces.** In this subsection, we construct open embeddings of a DFS $\pi : S \to X$ into projective $X$-schemes $\bar{\pi} : \bar{S} \to X$. Our procedure coincides with the one of Theorem 2 in [7] in case that $\pi : S \to X$ is defined over $\mathbb{C}$.

**4.6.** Given a DFS $\pi : S = S(\gamma) \to X$ defined by an essential $(A, x)$-labelled tree $\gamma$, we consider a factorization of the canonical morphism $\psi_\gamma : S = S_m \to \mathbb{A}^1_X = S_0$ into a sequence of fibered modifications $\beta_k : S_k \to S_{k-1}$, $k = 1, \ldots, m$. We embed $S_0$ in the $\mathbb{P}^1$-bundle $\bar{\pi}_0 : \bar{S}_0 := \mathbb{P}^1_X \to X$ as the complement a section $C \simeq X$ 'at infinity'. For every $k = 1, \ldots, m$, we let $\bar{\beta}_k : \bar{S}_k \to \bar{S}_{k-1}$ the blow-up morphism corresponding to the fibered modification $\beta_k : S_k \to S_{k-1}$. By construction, the morphism $\psi_\gamma$ lifts to an open embedding of $S$ into $\bar{\pi} : \bar{S} = \bar{S}_m \to X$. We denote by $E_j \simeq \mathbb{P}^1_k$, $j \in J$, the irreducible components of $\bar{\pi}^{-1}(F_0)$ different from the strict transform $F_0' \simeq \mathbb{P}^1_k$ of $F_0 = \bar{p}_0^{-1}(x_0)$ and the closures $\bar{C}_e \simeq \mathbb{P}^1_k$, $e \in L(\Gamma)$, of the irreducible components of $\pi^{-1}(x_0)$. Since for every $k = 1, \ldots, m$, the center of the blow-up $\bar{\beta}_k : \bar{S}_k \to \bar{S}_{k-1}$ is contained in the closure $\bar{D}_{k-1} \simeq \mathbb{P}^1_k$ in $\bar{S}_{k-1}$ of a unique irreducible component of $\bar{\beta}_{k-1}^{-1}(F_0) \cap S_{k-1}$, we conclude that $B_k = \bar{S}_k \setminus S_k$ is the strict transform of $B_{k-1} \cup \bar{D}_{k-1}$. Therefore, the dual graph of the $SNC$-divisor

$$\operatorname{Supp}\left(\bar{\pi}^{-1}(x_0)\right) = \overline{(B_m \setminus C')} \cup \bigcup_{e \in L(\Gamma)} \bar{C}_e$$

coincides with the underlying tree $\Gamma$ of $\gamma$. Since the tree $\Gamma$ is essential by assumption, we conclude that $\left({F'}_0^2\right) = -\mathrm{Card}\left(\mathrm{Child}_\Gamma(e_0)\right) \neq -1$, whereas $\left(E_j^2\right) \leq -2$ for every curve $E_j \subset \bar{S}$, $j \in J$. This leads to the following result.

**Proposition 4.7.** *A DFS $\pi : S = S(\gamma) \to X$ with base $(X, x_0)$ defined by a labelled tree $\gamma = (\Gamma, \sigma)$ admits an open embedding $i : S \hookrightarrow \bar{S}$ into a regular projective $X$-scheme $\bar{\pi} : \bar{S} \to X$ with the following properties.*

*a) $\bar{\pi}$ restricts to a trivial $\mathbb{P}^1$-bundle over $X_* = X \setminus \{x_0\}$.*

*b) The fiber $\bar{S}_{x_0} = \bar{\pi}^{-1}(x_0)$ is a reduced $SNC$-divisor which does not contain $(-1)$-curves meeting at most two other components transversally in a single point.*

*c) The irreducible components of $\bar{S}_{x_0}$ are isomorphic to $\mathbb{P}^1_k$.*

*d) The dual graph of $\bar{S}_{x_0}$ and $\bar{S}_{x_0} \setminus i(S_{x_0})$ are isomorphic to the underlying tree $\Gamma'$ of an essential tree for $\Gamma$ and to the subtree subtree $\Gamma' \setminus L(\Gamma')$ of $\Gamma'$ respectively.*

## 5. DANIELEWSKI SURFACES

We recall (2.1) that a *Danielewski surface is* a DFS with base $(k[x], x)$, where $k$ denotes a field of caracteristic zero. As a consequence of (1.6) and (3.2), the category $\overrightarrow{\mathfrak{D}}_{(k[x],x)}$ of relative Danielewski surfaces is equivalent to the category $\mathcal{T}^k_w$ of fine $k$-weighted trees. Therefore, every isomorphism class in $\overrightarrow{\mathfrak{D}}_{(k[x],x)}$ contains a canonical object $\left(\psi_0 : S_0 \to \mathbb{A}^1_X\right)$, consisting of the canonical morphism of the DFS $\pi : S_0 = S(\gamma_w) \to X$ defined by the $(k[x], x)$-labelled tree $\gamma_w$ obtain from a fine $k$-weighted tree by the procedure described in (1.6). In this section, we characterize Danielewski surfaces with a trivial Makar-Limanov invariant in case that the base field $k$ is algebraically closed.

*Notation* 5.1. Throughout this section, $\bar{k}$ denotes an algebraically closed field of caracteristic zero.

**Normal affine surfaces with a trivial Makar-Limanov invariant.** The Makar-Limanov invariant [10] of an affine algebraic variety $V = \mathrm{Spec}\,(B)$ over $\bar{k}$ is the sub-algebra $\mathrm{ML}\,(V)$ of $B$ consisting of regular functions on $V$ which are invariant under all $\mathbb{G}_{a,k}$-action on $V$. If $\mathrm{ML}\,(V) = \bar{k}$, then we say that $V$ has a trivial Makar-Limanov invariant. It is known that a normal affine surface surface $S$ has a trivial Makar-Limanov invariant if and only if it admits two $\mathbb{A}^1$-fibrations $\pi_1 : S \rightarrow X_1 \simeq \mathbb{A}^1_{\bar{k}}$ and $\pi_2 : S \rightarrow X_2 \simeq \mathbb{A}^1_{\bar{k}}$ with distinct general fibers. This property leads a geometrical criterion for $S$ to have trivial Makar-Limanov invariant. We recall that a minimal completion of $S$ is an open embedding $S \hookrightarrow \bar{S}$ of $S$ into a normal projective surface $\bar{S}$, nonsingular along $B = \bar{S} \setminus S$, such that $B$ is an $SNC$-divisor, with nonsingular curves as irreducible components, which does not contain $(-1)$-curves meeting at most two other components transversally in a single point. In [6], the author established that a complex normal affine surface $S$ has a trivial Makar-Limanov invariant if and only if the dual graph of the boundary divisor of any minimal completion of $S$ is a chain. Since the arguments given in Proposition 2.10 and Theorem 2.16 in [6] remain valid over an arbitrary algebraically closed field of caracteristic zero, we obtain the following more general criterion.

**Theorem 5.2.** *A normal affine surface* $S \not\simeq$ *Spec* $\left(\bar{k}\left[x, x^{-1}, y\right]\right)$ *has a trivial Makar-Limanov invariant if and only if the dual graph of the boundary divisor of every minimal completion of* $S$ *is a chain. Moreover, if there exists a completion of* $S$ *with this property, then this also holds for every other minimal completion.*

**Danielewski surfaces with a trivial Makar-Limanov invariant.** Here we apply the general criterion (5.2) to Danielewski surfaces.

**Definition 5.3.** A *comb* is a rooted tree $\Gamma$ such that $\Gamma \setminus L\,(\Gamma)$ is a chain. Equivalently, $\Gamma$ is a comb if every $e \in \Gamma \setminus L\,(\Gamma)$ has at most one child which is not a leaf of $\Gamma$. For instance, the tree $\Gamma$ of Figure 1 is a comb rooted in $e_0$.

For Danielewski surfaces

**Theorem 5.4.** *A Danielewski surface* $S/\bar{k}$ *has a trivial Makar-Limanov invariant if and only if it is isomorphic to a Danielewski surface* $S\,(\gamma)$ *defined by an* $\left(\bar{k}\,[x]\,, x\right)$*-labelled comb* $\gamma$.

*Proof.* A tree $\Gamma$ is a comb if and only if its essential subtree $\mathrm{Es}\,(\Gamma)$ is. Thus, by (3.10), we can suppose that $S = S\,(\gamma)$ for a certain essential $\left(\bar{k}\,[x]\,, x\right)$-labelled tree $\gamma = (\Gamma, \sigma)$. If $\gamma$ is the trivial tree, then $S \simeq \mathbb{A}^1_X \simeq \mathbb{A}^2_{\bar{k}}$ has trivial Makar-Limanov invariant. Otherwise we let $p_0 : \bar{S}_0 = \mathbb{P}^1_{\bar{X}} \rightarrow \bar{X}$, where $\bar{X} \simeq \mathbb{P}^1_X$ denotes a nonsingular projective model of $X \simeq \mathbb{A}^1_{\bar{k}}$, and we embed $S_0 = \mathbb{A}^1_X$ into $\bar{S}_0$ as the complement of the ample divisor $F_\infty \cup C$, where $F_\infty \simeq \mathbb{P}^1_{\bar{k}}$ denotes the fiber of $p_0$ over $\infty = \bar{X} \setminus X$, and where $C \simeq \mathbb{P}^1_{\bar{k}}$ is a section of $p_0$ 'at infinity'. By applying the same procedure as in (4.6) to $\bar{S}_0$, we obtain a projective surface $\bar{S}$ and an open embedding $S \hookrightarrow \bar{S}$ such that the dual graph $G\,(B)$ of the boundary divisor $B = \bar{S} \setminus S$ is isomorphic to the tree obtained from $\Gamma$ by deleting its leaves and replacing its root by a chain with three elements corresponding to the strict transforms $F_0', C'$ and $F_\infty'$ of the curves $F_0 = p_0^{-1}\,(x_0)$, $C$ and $F_\infty$. Since $\left(C'^2\right) = \left({F_\infty'}^2\right) = 0$ and $\left({F_0'}^2\right) \leq -2$ as $\gamma$ is essential, we conclude that $\bar{S}$ is a minimal completion of $S$. So the statement follows from (5.2) as $G\,(B)$ is a chain if and only if $\Gamma$ is a comb. $\square$

**Example 5.5.** Let $P_1, \ldots, P_n \in \bar{k}[T]$ be a collection of polynomials with simple roots, one of these roots, say $\lambda_{i,1} \in \bar{k}$, $1 \leq i \leq n$, being distinguished, and let

$$R_i(T) = (T - \lambda_{i,1})^{-1} P_i(T) = \prod_{j=2}^{r_i} (T - \lambda_{i,j}), \quad i = 1, \ldots, n.$$

The nonsingular surface $S = S_{P_1,\ldots,P_n} \subset \text{Spec}\left(\bar{k}[X_0, X_1, \ldots, X_{n+1}]\right)$ with equations

$$\left\{ \begin{array}{lcll} X_0 X_{j+1} & = & \prod_{i=1}^{j-1} R_i(X_i) P_j(X_j) & \text{for } 1 \leq j \leq n \\ (X_{j-1} - \lambda_{j-1,1}) X_{l+1} & = & X_j \prod_{i=j}^{l-1} R_i(X_i) P_l(X_l) & \text{for } 2 \leq j \leq l \leq n \end{array} \right.$$

where $\prod_{i=j}^{l-1} R_i(X_i) = 1$ if $j > l-1$, is a Danielewski surface with base $\left(\bar{k}[x], x\right)$. Indeed, $S\mid_{X_*} \simeq \text{Spec}\left(\bar{k}\left[X_0, X_0^{-1}\right][X_1]\right)$ whereas the irreducible components of the fiber $pr_x^{-1}(0)$ are the curves $C(\lambda_{1,1}, \ldots, \lambda_{k,1}, \ldots, \lambda_{m-1,j}) \simeq \mathbb{A}^1_{\bar{k}}$ with equations

$$\left\{ X_0 = 0, (X_i = \lambda_{i,1})_{i=1,\ldots,m-2}, X_{m-1} = \lambda_{m-1,j} \right\} \quad m = 1, \ldots, n,\ j = 1, \ldots, r_m$$

Moreover, the projection $p' : S \to \text{Spec}\left(\bar{k}[X_{n+1}]\right)$ is a second $\mathbb{A}^1$-fibration, and so $ML(S) = \bar{k}$. In [5], we prove that a Danielewski surface with base $\left(\bar{k}[x], x\right)$ has a trivial Makar-Limanov invariant if and only if it is isomorphic to a surface $S_{P_1,\ldots,P_n}$.

**Corollary 5.6.** *Given a morphism a Danielewski surfaces $\beta : S' \to S$ , we let*

$$\beta : S' = S_m \overset{\beta_m}{\to} S_{m-1} \overset{\beta_{m-1}}{\to} \cdots \overset{\beta_1}{\to} S_0 \overset{i}{\hookrightarrow} S$$

*be the factorization of $\beta$ into a sequence of fibered modifications followed by an open immersion. Then the following hold.*

*a) If $ML(S_{k_0}) = k$ then this also holds for $S_k$, $0 \leq k \leq k_0 - 1$.*

*b) If $ML(S) = k$ then this also holds for $S_0$.*

*Proof.* The Makar-Limanov invariant of $S_{k_0}$ being trivial, we can suppose that $S_{k_0}$ is isomorphic to the surface defined by a $\left(\bar{k}[x], x\right)$-labelled comb $\gamma = (\Gamma, \sigma)$. Then $S_{k_0-1}$ is isomorphic to the surface defined by a comb obtained from $\gamma$ by blowing-down the leaves at the unique maximal element $e$ of $\Gamma \backslash \text{Leaves}(\Gamma)$. Thus $S_{k_0-1}$ has a trivial Makar-Limanov invariant by virtue of (5.4), and so, (a) follows by induction. To prove (b), we can again suppose that $S \simeq S(\gamma)$ for a certain $\left(\bar{k}[x], x\right)$-labelled comb $\gamma = (\Gamma, \sigma)$. Then $S_0$ is isomorphic to the surface defined by the comb obtained from $\gamma$ by deleting a subset $N \subset \text{Leaves}(\Gamma)$. □

As a consequence of the above description, we recover the characterization of the Danielewski surfaces $S_{P,1} = \{xz - P(y) = 0\}$ in $\mathbb{A}^3_{\bar{k}} = \text{Spec}\left(\bar{k}[x, y, z]\right)$ given in [1].

**Corollary 5.7.** *For a Danielewski surface $\pi : S \to \mathbb{A}^1_{\bar{k}}$, the following are equivalent.*

*a) $ML(S) = \bar{k}$ and the canonical sheaf $\omega_S$ of $S$ is trivial*

*b) $S \simeq S(\gamma)$ for a suitable $\left(\bar{k}[x], x\right)$-labelled comb $\gamma$ of height $h \leq 1$.*

*c) $S \simeq S_{P,1} \subset \mathbb{A}^3_{\bar{k}}$ for a certain nonconstant polynomial $P$ with simple roots.*

*Proof.* By (3.10), we can suppose that $S = S(\gamma)$ for a certain essential $\left(\bar{k}[x], x\right)$-labelled tree $\gamma$. The canonical sheaf $\omega_{S(\gamma)}$ is trivial if and only if if and only if the invertible sheaf $\mathcal{L}_{n,-\mu(\gamma)}$ on $X(n)$ is trivial (see the proof of (2.13)). By construction, this is the case if and only if all the leaves of $\gamma$ are at the same level. On the other hand, by (5.2), $\mathrm{ML}(S(\gamma))$ is trivial if and only $\gamma$ is a comb. Thus (a) and (b) are equivalent since an essential comb with all its leaves at the same level is either the trivial tree or a nontrivial comb of height 1. Finally, (b) and (c) are equivalent by virtue of (3.5). $\square$

**Cyclic quotients of Danielewski surfaces.** A closed fiber $S_{z_0}$ of an $\mathbb{A}^1$-fibration $q : S \rightarrow Z$ on a normal affine surface is called *degenerate* if it is not isomorphic to $\mathbb{A}^1_{k(z_0)}$. In [7], Fieseler describes the structure of invariant neighbourhoods of degenerate fibers of a quotient $\mathbb{A}^1$-fibration $q : S \rightarrow Z = S//\mathbb{G}_{a,\mathbb{C}}$ on a normal affine surface $S/\mathbb{C}$ with a nontrivial $\mathbb{G}_{a,\mathbb{C}}$-action. This description, which remains valid for affine surfaces defined over an algebraically closed field of caracteristic zero, can be reinterpreted as follows.

**Proposition 5.8.** *Let $q : S \rightarrow Z$ be an $\mathbb{A}^1$-fibration on a normal affine surface $S/\bar{k}$. If $S_{z_0} = q^{-1}(z_0)$ is a degenerate fiber of $q$, then there exist a finite morphism $\phi : X \rightarrow Z' = Spec(\mathcal{O}_{Z,z_0})$ totally ramified at the unique point $x_0$ over $z_0$ and a DFS $\pi : \tilde{S} \rightarrow X$ with base $(X, x_0)$ equipped with an action of a cyclic group $\mathbb{Z}_m$ such that $S \times_Z Z' = \tilde{S}/\mathbb{Z}_m$.*

*Proof.* If $S_{z_0}$ is reduced then every irreducible component of $S_{z_0}$ is a connected component of $S_{z_0}$ and is isomorphic to $\mathbb{A}^1_{\bar{k}}$ (see Lemma 1.2 in [7]). Thus $p_2 : S' = S \times_Z Z' \rightarrow Z'$ is already a DFS with base $(Z', z_0)$. Otherwise, if $S_{z_0}$ is irreducible but not reduced, say with multiplicity $m \geq 2$, then, by Theorem 1.7 in [7], there exists a Galois covering $\phi : X \rightarrow Z'$ of order $m$, étale over $Z' \setminus \{z_0\}$ and totally ramified at the unique point $x_0$ over $z_0$, such that the normalization $\tilde{S}$ of $(S' \times_{Z'} X)_{\mathrm{red}}$ is a DFS $\pi : \tilde{S} \rightarrow X$, equipped with an action of the Galois group $\mathbb{Z}_m$ of the covering $\phi$ such that $S' \simeq \tilde{S}/\mathbb{Z}_m$. If $S_{z_0}$ is neither irreducible nor reduced, with irreducible components $C_i$ of multiplicity $m_i \geq 1$, $i = 1, \ldots, n$, then there exist DFS's $\pi_i : \tilde{S}_i \rightarrow X_i$ as above such that $S'_i = (S' \setminus S_{z_0}) \cup C_i$ is isomorphic to $\tilde{S}_i/\mathbb{Z}_{m_i}$ for every $i = 1, \ldots, n$. Letting $m$ be a common multiple of the $m_i$'s, these DFS's glue in a similar way as in the proof of Theorem 1.8 in [7] to a DFS $\pi : \tilde{S} \rightarrow Y$, equipped with an action of $\mathbb{Z}_m$ such that $S' \simeq \tilde{S}/\mathbb{Z}_m$. This completes the proof. $\square$

The above description is local in nature. However, a similar argument shows that a surface $S/\bar{k}$ equipped with an $\mathbb{A}^1$-fibration $q : S \rightarrow \mathbb{A}^1_{\bar{k}}$ with at most one degenerate fiber is isomorphic to cyclic quotient of a Danielewski surface. By Proposition 2.15 in [6], a normal affine surface $S/\bar{k}$ with a trivial Makar-Limanov admits an $\mathbb{A}^1$-fibration $q : S \rightarrow \mathbb{A}^1_{\bar{k}}$ with this property. So we obtain the following result.

**Theorem 5.9.** *A normal affine surface $S/\bar{k}$ with a trivial Makar-Limanov invariant is isomorphic to a cyclic quotient of a Danielewski surface.*

If the corresponding quotient morphism $\psi : S' \rightarrow S'/\mathbb{Z}_m \simeq S$ is étale in codimension 1, then the Danielewski surface $S'$ has a trivial Makar-Limanov invariant too (see *e.g.* [15]) . In general this is not the case, as shown by the following example.

**Example 5.10.** A similar argument as in (3.5) shows that the Danielewski surface with base $(\mathbb{C}[x], x)$ defined by the following labelled tree

$\gamma =$



is isomorphic to the surface $pr_x : S' \rightarrow \mathrm{Spec}\left(\mathbb{C}\left[x\right]\right)$ in $\mathbb{A}^4_{\mathbb{C}} = \mathrm{Spec}\left(\mathbb{C}\left[x,y,z,u\right]\right)$ given by the equations

$$xz - y\left(y^2-1\right) = 0, \quad \left(y^2-1\right)u - z^2 = 0, \quad xu - yz = 0.$$

It comes equipped a $\mathbb{Z}_2$-action $(x,y,z,u) \mapsto (-x,-y,z,u)$. The quotient $S/\mathbb{Z}_m$ is isomorphic to the nonsingular surface $S' \subset \mathrm{Spec}\left(\mathbb{C}\left[a,b,c,d,e\right]\right)$ with equations

$$\left\{ \begin{array}{c} ab = c^2, \quad ad = (b-1)\,c, \quad ae = cd = b\,(b-1)\,, \\ ce = bd, \quad (b-1)\,e = d^2 \end{array} \right. ,$$

via the quotient morphism $\phi : S' \rightarrow S$ induced by the ring homomorphism

$$\mathbb{C}\left[a,b,c,d,e\right] \quad \rightarrow \quad \mathbb{C}\left[x,y,z,u\right], (a,b,c,d,e) \mapsto \left(x^2, y^2, xy, z, u\right).$$

Note that $\phi$ is ramified along $pr_x^{-1}(0)$. Since $\gamma$ is not comb, $\mathrm{ML}\left(S'\right)$ is not trivial. On the other hand, $S$ has trivial Makar-Limanov invariant since it admits two $\mathbb{A}^1$-fibrations $pr_a : S \rightarrow \mathrm{Spec}\left(\mathbb{C}\left[a\right]\right)$ and $pr_e : S \rightarrow \mathrm{Spec}\left(\mathbb{C}\left[e\right]\right)$ with distinct general fibers.

## References

1. T. Bandman and L. Makar-Limanov, *Affine surfaces with* $AK(S) = \mathbb{C}$, Michigan J. Math. **49** (2001), 567–582.
2. N. Bourbaki, *Algèbre Commutative*, Éléments de Maths., vol. 27,28,30,31, Hermann, Paris, 1961-1965.
3. D. Daigle and P. Russell, *On log* $\mathbb{Q}$*-homology planes and weighted projective planes*, To appear in Canadian J. Math.
4. W. Danielewski, *On a cancellation problem and automorphism groups of affine algebraic varieties*, Preprint Warsaw, 1989.
5. A. Dubouloz, *Embeddings of generalized Danielewski surfaces in affine spaces*, (2003), E-print math.AG/0403208.
6. A. Dubouloz, *Completions of normal affine surfaces with a trivial Makar-Limanov invariant*, Michigan J. Math. **52** (2004), no. 2, 289–308.
7. K.H. Fieseler, *On complex affine surfaces with* $\mathbb{C}_+$*-actions*, Comment. Math. Helvetici **69** (1994), 5–27.
8. M.H. Gizatullin and V.I. Danilov, *Automorphismes of affine surfaces II*, Math. USSR Izvestiya **1** (1977), 51–98.
9. A. Grothendieck et al., *SGA1. Revêtements étales et Groupe Fondamental*, Lecture Notes in Math., vol. 224, Springer-Verlag, 1971.
10. S. Kaliman and L. Makar-Limanov, *On the Russel-Koras contractible threefolds*, J. Algebraic Geom. **6** (1997), 247–268.
11. S. Kaliman and M. Zaidenberg, *Affine modifications and affine hypersurfaces with a very transitive automorphism group*, Transformation Groups **4** (1999), 53–95.
12. L. Makar-Limanov, *On groups of automorphisms of a class of surfaces*, Israel J. Math. **69** (1990), 250–256.
13. L. Makar-Limanov, *On the group of automorphisms of a surface* $x^ny = p(z)$, Israel J. Math. **121** (2001), 113–123.
14. M. Miyanishi and K. Masuda, *The additive group actions on* $\mathbb{Q}$*-Homology planes*, Ann. Inst Fourier **53** (2003), no. 2, 429–464.
15. W. Vasconcelos, *Derivations of commutative noetherian rings*, Math. Z. **112** (1969), 229–233.

Adrien Dubouloz
Institut Fourier, Laboratoire de Mathématiques
UMR5582 (UJF-CNRS)
BP 74 St Martin d'Hères Cedex, FRANCE
*E-mail address*: `adrien.dubouloz@ujf-grenoble.fr`